

Screening of selected Malaysian plants against several food borne pathogen bacteria

¹Abd Aziz, S.M., ¹Low, C.N., ⁴Chai, L.C., ¹Abd Razak, S.S.N.,
¹Selamat, J., ¹Son, R., ³Sarker, M.Z.I. and ^{1,2*}Khatib, A.

¹Center of Excellence for Food Safety Research, Faculty of Food Science and Technology, University Putra Malaysia, 43400 Serdang, Selangor Darul Ehsan, Malaysia

²Laboratory of Natural Product, Institute of Bioscience, Universiti Putra Malaysia, 43400 Serdang, Selangor Darul Ehsan, Malaysia

³Department of Pharmaceutical Technology, Faculty of Pharmacy, International Islamic University Malaysia, Kuantan Campus, Jalan Sultan Ahmad Shah, Bandar Indera Mahkota, 25200, Kuantan, Pahang D/M, Malaysia

⁴Microbiology Unit, Institute of Biological Science, Faculty of Science, University of Malaya, 50603 Kuala Lumpur, Malaysia

Abstract: This study was conducted to evaluate antimicrobial properties of ethanolic extracts of the leaves of *Nephelium lappaceum*, *Curcuma longa*, *Cinnamomum cassia*, *Durio zibethinus*, *Vitex trifolia*, *Amaranthus tricolor*, *Syzygium samarangense* and *Manihot esculenta*. Antibacterial properties of the extracts were studied against fifteen strains of different gram positive and gram negative pathogenic bacteria, including *Bacillus cereus*, *Listeria monocytogenes*, *Staphylococcus aureus*, *Vibrio para*, and *Escherichia coli* using the agar disk diffusion method. Among the tested extracts, only *Amaranthus tricolor* exhibited specific inhibition of one of the tested bacteria; *Bacillus cereus*. Using the microdilution method, its minimum inhibitory concentration (MIC) value was determined to be 20 mg/mL.

Keywords: Malaysian plants, food borne pathogens, antimicrobial activity

Introduction

In an era of growing populations, the size of food chains increases and the challenge of producing safer food become increasingly tough. The official figures concerning food safety in Malaysia released by the Malaysian Ministry of Health (2009) revealed a sharp increase in reported food poisoning cases from 6930 to 17320 cases between 2006 and 2008. Moreover, in U.S, a country that is reknown for having safer food sources, the Center for Disease Control and Prevention estimated that 76 million people sicken, 300000 are hospitalized and 5000 people die each year between year 1991 and 2000 as a consequence of food poisoning (Olsen *et al.*, 2000).

The use of herbs to prevent and cure disease have been rising in recent years (López-Muñoz *et al.*, 2006) and its preventive use for the treatment of food borne pathogens is believed to be safer than the inclusion of synthetic antibiotics due to the long history of herbal usage. It is important, however, to consider issues such as the dosage and frequency of consumption of these products (Thomson *et al.*, 2001). Moreover, spices and herbs are able to enhance digestion, due to the salivary and gastric secretion induced by the pungent taste and unique aroma. This effect increase the activity of digestive enzymes, bile volume, and bile acids secretion, and also can decrease the time food

takes to travel through the gastrointestinal tract thus providing some protection against gastrointestinal disease (Low Dog, 2006).

In India, two herbs, *Vetvier* (*Vetiveria zizanoides*) and *Tulsi* (*Ocimum basilicum*), are commonly used for the treatment of food borne pathogens (Bais *et al.*, 2002; Champagnat *et al.*, 2008). Chinese use linseed, dandelion, golden thread and skullcap root possibly due to the presence of phenolic compounds (Pérez *et al.*, 1994; Lokar *et al.*, 1988; Franzblau *et al.*, 1986; Yu *et al.*, 2004), while ginseng and licorice have been used in Korea and Japan (Chang-Xiao *et al.*, 1992; Gupta *et al.*, 2008) for these purposes.

In Malaysia and Indonesia several native herbs have been traditionally used to treat food borne pathogens (Lu *et al.*, 2006; Arya *et al.*, 1989; Soedibyo, 1998; Chan *et al.*, 2007; Arora *et al.*, 1999; Aziz *et al.*, 2009; Ikram *et al.*, 2009). However, this use is lack of scientific proof and validation. In the third Herbal Asia Business Dialogue 2009, Kuala Lumpur, the prime goal was to transform Malaysian herbs production into the economy of tomorrow. To achieve that, incorporation of modern facilities to the production of traditional medicine has been widely developed.

The aim of this study was to evaluate the antimicrobial activity of the ethanol extracts of the leaves of *Nephelium lappaceum*, *Curcuma*

*Corresponding author.

Email: alfikhatib@hotmail.com

Tel: +6(03)89468540; Fax : +6(03)89423552

longa, *Cinnamomun cassia*, *Durio zibethinus*, *Vitex trifolia*, *Psidium guajava*, *Amaranthus tricolor*, *Syzygium samarangense* and *Manihot esculenta* against 5 pathogenic gram positive and negative bacteria: *Bacillus cereus*, *Listeria monocytogenes*, *Staphylococcus aureus*, *Vibrio para*, and *Escherichia coli*.

Materials and Methods

Chemicals

Tetracycline 30 µg antibiotics disc was purchased from Oxoid (Cambridge, UK), Nutrient agar, Mueller Hinton agar and broth, dimethylsulfoxide and ethanol were purchased from Merck (Darmstadt, Germany).

Herb selection and extraction

Herbs were selected based on the traditional usage of these plants for the treatment of food borne pathogen disease as listed in Table 1. Nine types of plants were collected from Botanical Farm Putrajaya in Malaysia on December 1, 2009 at 9.00-11.00 am. The herbs were identified by botanist, Mr. Mohammad Ishak. The voucher specimens were stored in Faculty of Food Science and Technology, Universiti Putra Malaysia. Dirt was removed under running water, and air dried for 3 days at constant room temperature 27°C with 70% relative humidity. Plants were then ground to powder using a mechanical grinder. A volume of 70 ml of ethanol was added to 10 g of plant powder, and the mixture was sonicated for 50 min at 30°C and 40 kHz frequency. The extracts were then filtered through filter paper and concentrated under reduced pressure using a rotary evaporator at 40°C for 15 min. Plant extracts were prepared by triplicate. The crude extracts were then kept at -20°C until used for analysis.

Table 1. List of selected plant used for study and related folk indication

No.	Scientific name	Malaysian local name	Part of the plant used	Folk indication
1	<i>Nephelium lappaceum</i>	<i>Daun rambutan</i>	leaf	Fever, diarrhea
2	<i>Vitex trifolia</i>	<i>Lemuni putih</i>	leaf	Diarrhea and gastrointestinal affections
3	<i>Curcuma longa</i>	<i>Kunyit</i>	leaf	Bloating
4	<i>Cinnamomun verum</i>	<i>Kayu manis</i>	leaf	Diarrhea and headache
5	<i>Syzygium samarangense</i>	<i>Jambu madu</i>	Shoot	Diarrhea
6	<i>Manihot esculenta</i>	<i>Ubi kayu</i>	leaf	Belly pain, diarrhea, fever, headache, aches and pains
7	<i>Psidium guajava</i>	<i>Jambu</i>	leaf	Headache, diarrhea
8	<i>Amaranthus tricolor</i>	<i>Bayam merah</i>	leaf	Sinus and cold fever, diarrhea
9	<i>Durio zibethinus</i>	<i>Durian</i>	leaf	Fever, diarrhea

Cited from Anderson *et al.* (1996); Arqueta *et al.* (1994); Chooi (2003); Chooi (2004); Soedibyo (1998)

Bacterial strains and inoculum preparation

Three strains of five types of pathogenic bacteria were selected for this study. The Gram positive bacteria were *Bacillus cereus* strains BC205, BC66, BC41, *Listeria monocytogenes* strains LM1, LM3, LM22 and *Staphylococcus aureus* strains SA25, SA2, SA5, while the gram negative bacteria were *Vibrio para* strains VB34, VB36, VB37, and *Escherichia coli* strains EC6, EC8, EC5. Growth media was Nutrient Agar from Merck (Darmstadt, Germany). Each bacteria was activated by transferring a loopful of culture onto the agar plate and after overnight incubation, all cultures were kept at -20°C in 50% glycerol stock to be used no longer than a month later.

Preliminary screening for antimicrobial activity

The disk diffusion agar method was applied, following National Committee for Clinical Standards (NCCLS, 2000). Mueller Hinton agar from Merck (Darmstadt, Germany) was used as the growth media of the pathogenic bacteria. After conformation with 0.5 Mc Farland turbidity, a sterile cotton swab was used to swab the culture evenly on the plate. The nine different extracts obtained were diluted in 10% DMSO to reach a concentration of 40 mg/mL. Sterile Whatman No.1 disk (6 mm) were soaked in the diluted extract for 1 min then dried completely before repeating the procedure until a total of 10 µL of the sample extract was absorbed onto the disk. Each extract was tested in triplicate with DMSO 10% as a negative control, and Tetracycline 30 µg was used as a positive control. Agar plates containing bacteria and extracts were incubated at 35°C for 24 hr. Size of inhibition ring was measured as the difference in the diameter between the growth-free zone and the disc.

Minimum inhibitory concentration

Sterile flat bottom tissue culture plates (96 well) obtained from Jet-Biofil® (Guangzhou, China) were filled with 100 µL Mueller Hinton broth (Darmstadt, Germany), 50 µL of the herbs extract and 50 µL bacterial suspension were inoculated into them. A stock solution of 40 mg/mL of herbs extract was prepared in DMSO 10%. An aliquot of this solution was serially diluted (two-fold) with nutrient broth to obtain a concentration range of 40 to 0.313 mg/mL. Each extract was assayed by triplicate. Controls were prepared using the same method as explained above but without adding the microbial culture. Plates were incubated at 35°C for 24 hr aerobically. A Benchmark Plus microplate spectrophotometer from Bio-Rad Laboratories Inc. (California, US) with Microplate Manager® software was used to read the absorbance

at 600 nm. The lowest concentration of the herbal extract to inhibit bacteria was considered as the MIC value of the tested extract. All experiments were conducted by triplicate for each type of bacteria and the values exhibited are Mean \pm SD.

Results and discussions

There was a large variability in the antimicrobial activities displayed by the tested herbs during this screening as shown in Table 2. The classification of the activity as weak, medium, strong and very strong followed that of used by US National Committee for Clinical Laboratory Standards (2000) and Lin *et al.* (1999). About 44 % of herbs showed some degree of inhibition of both gram negative and/or gram positive bacteria. Among the active herbs, 57% inhibited gram positive bacteria and only 43% inhibited gram negative bacteria. Therefore, this proved that Gram negative bacteria were more resistant than gram positive, a fact explained by the presence of lipopolysaccharide in their outer wall (Russel, 1991).

In most previous research of this type, when an inhibitory effect of pathogenic bacteria growth was detected, the plant extracts responsible for this activity were reported to be capable of inhibiting various types of microorganisms. This result is considered to be a 'false-positive' result since the mechanism of inhibition could well be due to cell toxicity (Cos *et al.*, 2006). A plant that is able to inhibit only specific bacteria is the plant which is worthwhile studying. Out of the 9 herbs tested in this study, only *A. tricolor* showed a specific inhibition of one type of bacteria, *B. cereus*. This is the first report on the specific antimicrobial activity of *A. tricolor* ethanolic extract against *B. cereus*. Sharma (1993) reported non-specific antimicrobial activity of this plant extracted with petroleum ether on both gram positive and negative bacteria. The difference with the results obtained in this study is very likely due to difference in polarity of the two solvents used which would result in different types of compounds being extracted. As shown in Figure 1, the MIC values of the *A. tricolor* extracts calculated for all strains of *B. cereus* was 20 mg/mL. A MIC value below 50 mg/ml is considered to correspond to an interesting antibacterial activity which is worth while continuing to study (Arias *et al.*, 2004; Motamedi *et al.*, 2010; Safary *et al.*, 2009; Fabry *et al.*, 1998; Valsaraj *et al.*, 1997).

A. tricolor belongs to the Amaranthaceae family. Screening of antimicrobial activity on Amaranthaceae family members such as the ethanolic extract of aerial parts and roots of *Blutaparon portulacoides* showed




Figure 1. Minimum inhibitory concentration (MIC) of *A. tricolor* toward *B. cereus* strains

specific inhibition towards a gram positive bacteria, *Streptococcus* sp (Salvador *et al.*, 2002). Another study on plants from this family revealed that an 80% ethanol extract of leaf and stems of *Achyranthes aspera*, also had a specific inhibition towards two gram positive bacteria, *B. subtilis* and *S. aureus* (Valsaraj *et al.*, 1997) while the methanolic and water extract of leaves of *Alternanthera brasiliana*, were reported to present inhibition to both gram positive and gram negative bacteria at 25 mg/ml (Coelho de Souza *et al.*, 2004).

Several sterols such as cholesterol, 7,22-ergostadienol, campesterol, stigmasterol, 7,24(28)-ergostadienol, 7-ergostenol, spinasterol, sitosterol, stigmastanol, 7, 25-stigmastadienol, 7-stigmastenol, and 7,24(28)-stigmastadienol have been isolated from *A. tricolor* (Sharma, 1993; Xu *et al.*, 1986). Alkanol groups that were found to be present were hexacosanol, and octacosanol while the alkanes nonacosane, and hentriaconetane were also reported (Sharma, 1993). The presence of β -sitosterol and stigmasterol in *A. tricolor* has been related to a strong inhibition against gram-positive bacteria and a moderate activity against gram negative bacteria (Sharma, 1993). Alkanes are the most inert organic compound to give wide spectrum of antibacterial effect (Sharma, 1993). However, no compounds from *A. tricolor* had been reported to have specific inhibition towards *B. cereus*.

The other ethanolic plant extracts tested showed non-specific antibacterial activity. This was the case of *N. lappaceum*, *S. samarangense*, and *V. trifolia* extracts. The ethanolic extract of *N. lappaceum* leaves strongly inhibited one strain of *S. aureus* and *V. para*, and showed medium inhibition towards one strain of *L. monocytogenes*. The other strains of *S. aureus* were weakly inhibited. From the previous literature, only the rind of this plant, extracted with ether, methanol, and water, showed potent activity against several gram positive bacteria including *S. aureus*, *S. epidermis*, and *E. faecalis* and only one gram negative bacteria, *V. cholera*. Phenolic compounds from the peel of this plant were related to

Table 2. Antimicrobial activity of leaf extracts on selected bacterial strains^a

No	Common name	Gram positive bacteria									Gram negative bacteria					
		<i>Bacillus cereus</i>			<i>Listeria monocytogenes</i>			<i>Staphylococcus aureus</i>			<i>Vibrio para</i>			<i>Escherichia coli</i>		
		BC205	BC66	BC41	LM1	LM3	LM22	SA25	SA2	SA5	VP37	VP36	VP34	EC6	EC8	EC5
1	<i>N. lappaceum</i>	-	-	-	-	-	++	+++	+	+	+++	-	-	-	-	-
2	<i>V. trifolia</i>	+++	+++	-	+++	++	+++	++	++++	++	-	-	-	+++	-	++++
3	<i>C. longa</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4	<i>C. verum</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5	<i>S. samarangense</i>	-	-	-	-	-	+	-	-	-	+	-	-	++	++	++
6	<i>M. esculenta</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
7	<i>P. guajava</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
8	<i>A. tricolor</i>	++++	++++	+++	-	-	-	-	-	-	-	-	-	-	-	-
9	<i>D. zibethinus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10	DMSO	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
11	Te	+++	++++	++++	++++	+++	++++	-	-	-	+++	++++	++++	++++	++++	++++

^a+: 2-4 mm, weak inhibition; ++: 5-7 mm, medium inhibition; +++: 8-10 mm, strong inhibition; ++++: >10 mm, very strong inhibition, excluding the diameter of the disc; -: no activity; Te: Tetracycline 30µg/disc.

this antibacterial activity (Thitilertdecha *et al.*, 2008) and more antibacterial activity was found in the peel than in the seed. However, no antimicrobial study had been conducted on the leaves of this plant.

Ethanol extract of *S. samarangense* leaves showed medium inhibition towards all strains of *E. coli*, and weak inhibition towards one strain of *L. monocytogenes* and *V. para*. Previous phytochemical studies on the leaves of *S. samarangense* showed the presence of ellagitannins, proanthocyanidins (Nonaka *et al.*, 1992), flavanones (Kuo *et al.*, 2004), flavonol glycosides (Kuo *et al.*, 2004; Nair *et al.*, 1999), anthocyanidins (Nonaka *et al.*, 1992; Kuo *et al.*, 2004), triterpenoids (Srivastaya *et al.*, 1995), chalcones (Srivastaya *et al.*, 1995; Resurreccion-Magno *et al.*, 2005), and volatile terpenoids (Wong *et al.*, 1996). Ellagitannin had been reported to have a specific antimicrobial activity against *S. aureus* (Perashar *et al.*, 2009). Triterpenoids showed inhibition towards both *B. subtilis* and *E. coli* (Nick *et al.*, 1994) while flavonoids exhibited antibacterial activity to both gram negative and gram positive (Cushnie *et al.*, 2005; Cook *et al.*, 1996).

The ethanol extract of *V. trifolia* exhibited an inhibitory effect against all tested bacteria of strong to medium strength with the exception of *V. para*. This finding is in accordance with the result reported by Hernández *et al.* (1999) who used hexane, dichloromethane and methanolic extracts of this plant to inhibit a wide range of bacteria such as *S. aureus*, *S. faecalis*, *E. coli*, *Proteus mirabilis*, *Shigella sonnei*, *Salmonella typhimurium*, and *Candida albicans*. The acetone extract of this plant showed inhibition towards gram negative and gram positive bacteria (Nyiligira *et al.*, 2008). Different groups of compounds had been isolated from this plant, including flavanoids such as casticin, vitexin, artemetin, luteolin (Zeng *et*

al., 1996), sterols, such as β -sitosterol (Nair *et al.*, 1975), terpenoids, monoterpenes, labdane diterpene and sesquiterpenes. (Nyiligira *et al.*, 2008; Hossain *et al.*, 2001; Tandon *et al.*, 2008; Singh *et al.*, 1999) These compounds have all been reported to have antibacterial activity (Cushnie *et al.*, 2005).

The ethanol extract of *P. guajava* and *C. cassia* leaves showed no inhibition towards the tested bacteria. This result coincided with the findings of previous studies which showed no antimicrobial activity of ethanol *P. guajava* leaf extract towards *S. aureus* and β -Streptococcus (Jaiarj *et al.*, 1999). The activity was different when other solvents were used for extraction as reported by Jaiarj (1999), Chang *et al.* (2001) and Wang *et al.* (2009). The water extract from *P. guajava* leaves showed high inhibitory effect towards *S. aureus* (Jaiarj *et al.*, 1999). Meanwhile, Chang (2001) and Wang (2009) reported that essential oils from *C. cassia* extracted by ethyl ether exhibited antibacterial activity.

Ethanol extract of *D. zibethinus*, *M. esculenta*, and *C. longa* leaves exhibited no inhibitory effect on the tested bacteria. No previous report has been found on the antimicrobial activity of these plants.

Conclusions

Some of the plant extracts tested in this study showed inhibition towards both gram negative and/or gram positive bacteria; however the only plant which exhibited a specific inhibition to certain bacteria was considered as the potential candidate for an antimicrobial agent. This study demonstrated that the ethanol extract of *A. tricolor* leaves specifically inhibited all tested strains of *B. cereus* with a MIC value of 20 mg/ml. *B. cereus* is normally found in products containing starch, meat, and vegetable. Some

B. cereus outbreaks have also occurred in “meal-on wheel”, where the food has been kept above room temperature for a period of time before being served to consumers. Some cases may originate in stir-fry products since the toxin produced by *B. cereus* is stable to heat (Granum *et al.*, 1997).

Acknowledgements

The authors wish to thank University Putra Malaysia for financial support through Research University Grant Scheme (RUGS) Initiative 4, and deputy of Putrajaya Botanical Farm, Noorsiha Ayub for supplying the plant materials. Erica G. Wilson is also acknowledged for English editing.

References

- Anderson, G. J. and Coee, F. G. 1996. Ethnobotany of the garifuna of East Nicaragua. *Economic Botany* 50: 71-107.
- Argueta, A., Cano, L. M., and Rodarte, M. E. 1994. Atlas de las Plantas de la Medicina Tradicional I y III, 537-538. Mexico.
- Arias, M. E., Gomez, J. D., Cudmani, N. M., Vattuone, M. A., and Isla, M. I. 2004. Antibacterial activity of ethanolic and aqueous extracts of *Acacia aroma* Gill. *ex Hook et Arn. Life Sciences* 75: 191-202.
- Arora, D. S. and Kaur, J. 1999. Antimicrobial activity of spices. *International Journal of Antimicrobial Agents* 12: 257-262.
- Arya, R., Babu, V., Ilyas, M., and Nassim, K. T. 1989. Phytochemical examination of the leaves of *Anacardium occidentale*. *Journal of the Indian Chemical Society* 66: 67-68.
- Aziz, Z. and Tey, N. P. 2009. Herbal medicines: Prevalence and predictors of use among Malaysian adults. *Complementary Therapies in Medicine* 17: 44-50.
- Bais, H. P., Walker, T. S., Schweizer, H. P., and Vivanco, J. M. 2002. Root specific elicitation and antimicrobial activity of Rosmarinic acid in hairy root cultures of *Ocimum basilicum*. *Plant Physiology and Biochemistry* 40: 983-995.
- Champagnat, P., Heitz, A., Carnat, A., Fraise, D., Carnat, A.-P., and Lamaison, J.-L. 2008. Flavonoids from *Vetiveria zizanioides* and *Vetiveria nigriflora* (Poaceae). *Biochemical Systematics and Ecology* 36: 68-70.
- Chan, E. W. C., Lim, Y. Y., and Omar, M. 2007. Antioxidant and antibacterial activity of leaves of *Etilingera* species (Zingiberaceae) in Peninsular Malaysia. *Food Chemistry* 104: 1586-1593.
- Chang, S.-T., Chen, P.-F., and Chang, S.-C. 2001. Antibacterial activity of leaf essential oils and their constituents from *Cinnamomum osmophloeum*. *Journal of Ethnopharmacology* 77: 123-127.
- Chang-Xiao, L. and Pei-Gen, X. 1992. Recent advances on ginseng research in China. *Journal of Ethnopharmacology* 36: 27-38.
- Chooi, O. H. 2003. Sayuran: Khasiat makanan dan ubatan, Utusan Publication and Distributors Sdn Bhd, Kuala Lumpur.
- Chooi, O. H. 2004. Tumbuhan liar, Khasiat ubatan dan kegunaan lain, Utusan Publications and Distributors Sdn Bhd.
- Coelho de Souza, G., Haas, A. P. S., von Poser, G. L., Schapoval, E. E. S., and Elisabetsky, E. 2004. Ethnopharmacological studies of antimicrobial remedies in the south of Brazil. *Journal of Ethnopharmacology* 90: 135-143.
- Cook, N. C. and Samman, S. 1996. Flavonoids-chemistry, metabolism, cardioprotective effects and dietary sources. *Journal of Nutritional Biochemistry* 7: 66-76.
- Cos, P., Vlietinck, A. J., Berghe, D. V., and Maes, L. 2006. Anti-infective potential of natural products: How to develop a stronger *in vitro* 'proof-of-concept'. *Journal of Ethnopharmacology* 106: 290-302.
- Cushnie, T. P. T. and J.Lamb, A. 2005. Antimicrobial activity of flavonoids. *International Journal of Antimicrobial Agents* 26: 343-356.
- Fabry, W., Okemo, P. O., and Ansorg, R. 1998. Antibacterial activity of East African medicinal plants. *Journal of Ethnopharmacology* 60: 79-84.
- Franzblau, S. G. and Cross, C. 1986. Comparative *in vitro* antimicrobial activity of chinese medicinal herbs. *Journal of Ethnopharmacology* 15: 279-288.
- Granum, P. E. and Lund, T. 1997. *Bacillus cereus* and its food poisoning toxins. *FEMS Microbiology Letters* 157: 223-228.
- Gupta, V. K., Fatima, A., Faridi, U., Negi, A. S., Shanker, K., Kumar, J. K., Rahuja, N., Luqman, S., Sisodia, B. S., Saikia, D., Darokar, M. P., and Khanuja, S. P. S. 2008. Antimicrobial potential of *Glycyrrhiza glabra* roots. *Journal of Ethnopharmacology* 116: 377-380.
- Hernández, M. M., Heraso, C., Villarreal, M. L., Vargas-Arispuro, I., and Aranda, E. 1999. Biological activities of crude plant extracts from *Vitex trifolia* L. (Verbenaceae). *Journal of Ethnopharmacology* 67: 37-44.
- Hossain, M. M., Paul, N., Sohrab, M. H., Rahman, E., and Rashid, M. A. 2001. Antibacterial activity of *Vitex trifolia*. *Fitoterapia* 72: 695-697.
- Ikram, E. H. K., Eng, K. H., Jalil, A. M. M., Ismail, A., Idris, S., Azlan, A., Nazri, H. S. M., Diton, N. A. M., and Mokhtar, R. A. M. 2009. Antioxidant capacity and total phenolic content of Malaysia underutilized fruits. *Food Composition and Analysis Journal* 22: 388-393.
- Jaiarj, P., Khoohaswan, P., Wongkrajang, Y., Peungvicha, P., Suriyawong, P., Saraya, M. L. S., and Ruangsomboon, O. 1999. Anticough and antimicrobial activities of *Psidium guajava* Linn. leaf extract. *Journal of Ethnopharmacology* 67: 203-212.
- Kuo, Y.-C., Yang, L.-M., and Lin, L.-C. 2004. Isolation and immunomodulatory effects of flavonoids from *Syzygium samarangense*. *Planta Medical Research Reviews* 70: 1237-1239.
- Lin, J., Opoku, A.R., Geheeb-Keller, M., Hutchings, A.D., Terblanche, S.E., Jäger, A.K., and van Staden, J.

- J. 1999. Preliminary screening of some traditional zulu medicinal plants for anti-inflammatory and antimicrobial activities. *Journal of Ethnopharmacology* 68: 267 - 274.
- Lokar, L. C. and Poldini, L. 1988. Herbal remedies in the traditional medicine of the Venezia Giulia Region (North East Italy). *Journal of Ethnopharmacology* 22: 231-279.
- López-Muñoz, F., Alamo, C., and García-García, P. 2006. "The herbs that have the property of healing...": The phytotherapy in Don Quixote. *Journal of Ethnopharmacology* 106: 429-441.
- Low Dog, T. 2006. A reason to season: The therapeutic benefits of spices and culinary Herbs. *The Journal of Science and Healing* 2: 446-449.
- Lu, H., Liang, Y., Yi, L., and Wu, X. 2006. Anti-inflammatory effect of *Houttuynia cordata* injection. *Journal of Ethnopharmacology* 104: 245 - 249.
- Malaysian Ministry of Health. 2009. Health Fact 2009, Health Informatics Center Planning and Development Ministry of Health Malaysia, Kuala Lumpur.
- Motamedi, H., Darabpour, E., Gholipour, M., and Nejad, S. M. S. 2010. Antibacterial effect of ethanolic and methanolic extracts of *Plantago ovata* and *Olivaria decumbens* endemic in Iran against some pathogenic bacteria. *International Journal of Pharmacology* 6: 117-122.
- Nair, A. G. R., Krishnan, S., Rayikrishnan, C., and Madhusudnan, K. P. 1999. New and rare flavonol glycosides from leaves of *Syzygium samarangense*. *Fitoterapia* 70: 148-151.
- Nair, A. G. R., Ramesh, P., and Subramanian, S. S. 1975. Two unusual flavones (artemetin and 7-desmethyl artemetin) from the leaves of *Vitex trifolia*. *Current Science* 44: 214-216.
- National Committee for Clinical Laboratory Standards. 2000. Performance standards for antimicrobials disk susceptibility tests. Approved standard 7th ed., NCCLS document M2-A7, Wayne, PA
- Nick, A., Wright, A. D., Sticher, O., and Rali, T. 1994. Antibacterial triterpenoid acids from *Dillenia papuana*. *Journal of Natural Products* 57: 1245-1250.
- Nonaka, G., Aiko, Y., Aritake, K., and Nishioka, I. 1992. Tannins and related compounds. CXIX Samarangenins a and b, novel proanthocyanidins with doubly bonded structures from *Syzygium samarangen* and *S. aquem*. *Chemical and Pharmaceutical Bulletin* 40: 2671-2673.
- Nyiligira, E., Viljoen, A. M., Van Heerden, F. R., Van Zyl, R. L., Van Vuuren, S. F., and Steenkamp, P. A. 2008. Phytochemistry and in vitro pharmacological activities of South African *Vitex* (Verbenaceae) species. *Journal of Ethnopharmacology* 119: 680-685.
- Olsen, S. J., MacKinnon, L. C., Goulding, J. S., Bean, N. H., and Slutsker, L. 2000. Centers for Disease Control and Prevention, MMWR 2000, March 17
- Perashar, A., Gupta, C., Gupta, S. K., and Kumar, A. 2009. Antimicrobial ellagitannin from pomegranate (*Punica granatum*) fruits's. *International Journal of Fruit Science* 9: 226-231.
- Pérez, C. and Anesini, C. 1994. In vitro antibacterial activity of Argentine folk medicinal plants against *Salmonella typhi*. *Journal of Ethnopharmacology* 44: 41-46.
- Resurreccion-Magno, M. H. C., Villasenor, I. M., Harada, N., and Monde, K. 2005. Antihyperglycemic flavonoids from *Syzygium samarangense* (Blume) merr and perry. *Phytotherapy Research* 19: 246-253.
- Russel, A. D. 1991. Mechanisms of bacterial resistance to non-antibiotics: food additives and food pharmaceutical preservatives. *Journal of Applied Bacteriology* 71: 191-201.
- Safary, A., Motamedi, H., Maleki, S., and Seyyednejad, S. M. 2009. A preliminary study on the antibacterial activity of *Quercus brantii* against bacterial pathogens, particularly enteric pathogens. *International Journal of Botany* 5: 176-180.
- Salvador, M. J., Ferreira, E. O., Pral, E. M. F., Alfieri, S. C., Albuquerque, S., Ito, I. Y., and Dias, D. A. 2002. Bioactivity of crude extracts and some constituents of *Blutaparon portulacoides* (Amaranthaceae). *Phytomedicine* 9: 566-571.
- Sharma, R. K. 1993. Phytosterols: Wide-spectrum antibacterial agents. *Bioorganic Chemistry* 21: 49-60.
- Singh, M., Pal, M., and Sharma, R. P. 1999. Biological activity of the labdane diterpenes. *Plant Medica* 68: 72-73.
- Soedibyo, M. 1998. Alam Sumber Kesehatan Manfaat dan Kegunaan, Balai Pustaka, Jakarta.
- Srivastaya, R., Shaw, A. K., and Kulshreshta, D. K. 1995. Triterpenoids and chalcone from *Syzygium samarangense*. *Phytochemistry* 38: 687-689.
- Tandon, S., Mittal, A. K., and Pant, A. K. 2008. Insect growth regulatory activity of *Vitex trifolia* and *Vitex agnus-castus* essential oils against *Spilosoma obliqua*. *Fitoterapia* 79: 283-286.
- Thitilertdech, N., Teerawutgulrag, A., and Rakariyatham, N. 2008. Antioxidant and antibacterial activities of *Nephelium lappaceum* L. extracts. *Food Science and Technology* 41: 2029-2035.
- Thomson, C., Lutz, R. B., Ann, M. C., Cheryl, L. R., and Elaine, R. M. 2001. Herbs and botanical supplements: Principles and concepts, Nutrition in the Prevention and Treatment of Disease, Academic Press, San Diego, pp. 261-275.
- Valsaraj, R., Pushpagandan, P., and Smitt, U. W., Adersen, A., Nyman, U. 1997. Antimicrobial screening of selected medicinal plants from India. *Journal of Ethnopharmacology* 58: 75-83.
- Wang, R., Wang, R., and Yang, B. 2009. Extraction of essential oils from five cinnamon leaves and identification of their volatile compound compositions. *Innovative Food Science & Emerging Technologies* 10: 289-292.
- Wong, K. C. and Lai, F. Y. 1996. Volatile constituents from the fruits of four species grown in Malaysia. *Journal of Flavour and Fragrance* 11: 61-66.
- Xu, S., Patterson, G. W., and Schmid, K. 1986. Sterols of

amaranthaceae. *Phytochemistry* 25: 1883-1886.

- Yu, J., Lei, J., Yu, H., Cai, X., and Zou, G. 2004. Chemical composition and antimicrobial activity of the essential oil of *Scutellaria barbata*. *Phytochemistry* 65: 881-884.
- Zeng, X., Fang, Z., Wu, Y., and Zhang, H. 1996. Chemical constituents of the fruits of *Vitex trifolia* L. . *Chung-Kuo Chung Yao Tsa Chih* 21: 167-168.