

## Mini Review

## Flaxseed – a nutritional punch

\*Ganorkar, P. M. and Jain, R. K.

Department of Food Processing Technology,  
A.D. Patel Institute of Technology, New Vallabh Vidya Nagar, Anand,  
Gujarat 388121, India

### Article history

Received: 27 June 2012  
Received in revised form:  
7 September 2012  
Accepted: 5 October 2012

### Abstract

Flaxseed is mainly considered as oilseed crop. Moreover, the other nutritional parameters than its oil content, make it more favorable choice for food technologist to develop functional foods. Flaxseed contains good amount of  $\alpha$ -Linolenic Acid (ALA), a omega-3 fatty acid, protein, dietary fiber, lignan, specifically Secoisolariciresinol diglucoside (SDG). Several studies reveal that these components work well for nutritional benefit in human being. ALA is beneficial for infant brain development, reducing blood lipids and cardiovascular diseases. Flaxseed proteins are relatively high in arginine, aspartic acid and glutamic acid whereas lysine, methionine and cystine are limiting amino acid. Flaxseed dietary fiber exhibits positive effect to reduce constipation, to keep better bowel movement and as hypocholestermic agent. SDG have antioxidant activity and free oxygen radical scavenging activity. Consequently, it may have anticancer property. At some extent, SDG helps in bone development. Cyanogenic glycosides and linatine are antinutrients in flaxseed. As compared to soyabean and canola, flaxseed antinutrient effect on human health is very less. Researchers reported that flaxseed incorporated food products can have good consumer acceptability along with its nutritional benefits.

### Keywords

ALA  
protein  
dietary fiber  
lignan  
cyanogenic glucosides  
linatine

© All Rights Reserved

### Introduction

Flaxseed, or Linseed (*Linum Usitatissimum*), popularly known as Alsi, Jawas, Aksebija in Indian languages, is a blue flowering rabi crop and a member of family Linaceae. Annual production of flax was 3.06 million tons and Canada is the world's largest producer of flax (about 38% of total production) (Anonymous, 2000). Globally, Flaxseed is grown as either oil crop or a fiber crop with fiber linen derived from the stem of fiber varieties and oil from the seed of linseed varieties (Diederichsen *et al.*, 2003; Vaisey-Genser *et al.*, 2003). The plant is native to west Asia and the Mediterranean. As the source of linen fiber flax has been cultivated since at least 5000 BC, today it is mainly grown for its oil (Berugland, 2002; Oomah, 2001). The spherical fruit capsules contain two seeds in each of five compartments. The seed is flat and oval with a pointed tip. It have smooth glossy surface. It varies in color dark brown to yellow (Freeman, 1995). The texture of flaxseed is crisp and chewy possessing a pleasant nutty taste (Carter, 1996). Beyond its oilseed crop ability, proximate

composition of flaxseed makes it more promising for its utilization in different food products. Flaxseed is one of the richest vegetarian source of  $\alpha$ -linolenic acid (omega 3 fatty acid) and soluble mucilage. In present era, consumer's trend towards functional food has increased significantly as health awareness rose. Flaxseed can be one stop for novel high quality source of nutrition.

### Proximate composition of flaxseed

Flax is rich in fat, protein and dietary fibre. An analysis of brown Canadian flax averaged 41% fat, 20% protein, 28% total dietary fibre, 7.7% moisture and 3.4% ash, which is the mineral-rich residue left after samples are burned (Morris, 2003). The composition of flaxseed can vary with genetics, growing environment, seed processing and method of analysis (Daun *et al.*, 2003). The protein content of the seed decreases as the oil content increases (Daun and DeClercq, 1994). The oil content of flaxseed can be altered through traditional plant breeding methods, and it is affected by geography – the cool nights of northern Canada improve oil content and quality. The

\*Corresponding author.  
Email: [pmganorkar@rediffmail.com](mailto:pmganorkar@rediffmail.com)

composition of flaxseed is shown in Table 1.

Brown and yellow (Omega) varieties of flaxseed are virtually identical in their nutrient content (Morris, 2003). Seed coat colour is determined by the amount of pigment present, a feature that can be changed through normal plant breeding practices. Consumers can buy brown or yellow flaxseed based on price and appearance of the flaxseed containing food product, since the nutritional value of brown and yellow flax is similar. Flaxseed oil and canola oil have the lowest levels of the nutritionally undesirable saturated fatty acids. The level of the desirable monounsaturates in flax oil is modest.

#### *Flaxseed as a source of ALA (Omega-3 fatty acid)*

There are two groups of omega fats: omega-3 and omega-6 fatty acids. Linolenic acid, eicosapentaenoic acid (EPA) and docosahexanoic acid (DHA) are three types of omega-3 fatty acids and are nutritionally important. All three fatty acids have been shown to reduce the risk of cardiovascular disease (Hurteau, 2004). Flax contains a mixture of fatty acids. It is rich in polyunsaturated fatty acids, particularly ALA, the essential omega-3 fatty acid, and linoleic acid (LA), the essential omega-6 fatty acid. These two polyunsaturated fatty acids are essential for humans – that is, the body needs them. Supercritical CO<sub>2</sub> extraction gave a higher average ALA content (60.5%) compared to the soxhlet extraction method (56.7%) (Bozan and Temelli, 2002). Fatty acid content of flaxseed oil is depicted in Table 2.

ALA and Linoleic acid constitutes 57% and 16.0 % of total fatty acids respectively in flax making the richest source of ALA. ALA from flaxseed exerts positive effect on blood lipids. It was found to be as effective as oleic acid (18:2 $\eta$ -6) and linoleic acid (18:2 $\eta$ -6) in the reduction of plasma total cholesterol, low density lipoprotein cholesterol and very low density lipoprotein cholesterol in 20-34 years old healthy men (Chan *et al.*, 1993). 12 g of ALA was taken three times a day by group of healthy young women in the flaxseed oil capsules and compared with group given in flaxseed flour supplemented products. Impressive reductions in blood lipids were observed in both cases (Cunnane *et al.*, 1993). Nettleton (2003) summarized the recommendations of leading health organizations regarding the proper ratio of n-6 to n-3 fatty acid intake. Most organizations agree that a 5:1 to 10:1 n-6 to n-3 fatty acid ratio is preferred (Institute of Medicine, 2002; WHO/FAO, 2003). However, a typical diet has an n-6 to n-3 fatty acid ratio well beyond 10:1; thus, flaxseed can be a valuable lipid source to improve the n-6 to n-3 fatty acid ratio due to the high n-3 content of flaxseed

Table 1. Proximate composition of flaxseed on common measures<sup>a</sup>

Form of flax	Weight(g)	Common measure	Energy (kcal)	Total fat (g)	ALA <sup>b</sup> (g)	Protein (g)	Total CHO <sup>c,d</sup> (g)	Total dietary fibre (g)
Proximate analysis	100	-	450	41.0	23.0	20.0	29.0	28.0
Whole seed	180	1 cup	810	74.0	41.0	36.0	52.0	50.0
	11	1 tbsp	50	4.5	2.5	2.2	3.0	3.0
	4	1 tsp	18	1.6	0.9	0.8	1.2	1.1
Ground seed	130	1 cup	585	53.0	30.0	26.0	38.0	36.0
	8	1 tbsp	36	3.3	1.8	1.6	2.3	2.2
	2.7	1 tsp	12	1.1	0.6	0.5	0.8	0.8
Flax oil	100	-	884	100.0	57.0	-	-	-
	14	1 tbsp	124	14.0	8.0	-	-	-
	5	1 tsp	44	5.0	2.8	-	-	-

<sup>a</sup>Based on a proximate analysis conducted by the Canadian Grain Commission. The fat content was determined using the American Oil Chemists' Society (AOCS) Official Method Am 2-93. The moisture content was 7.7%.

<sup>b</sup>ALA = Alpha-linolenic acid

<sup>c</sup>CHO = Carbohydrate.

<sup>d</sup>Total Carbohydrate includes carbohydrates like sugars and starches (1 g) and total dietary fibre (28 g) per 100 g flax seeds.

(Source: Morris, D. H. 2003. Flax: A health and nutrition primer. p. 12. Winnipeg: Flax Council of Canada)

Table 2. Fatty acids content of Flaxseed oil

Parameter	Percentage (%)
Saturated fat	9.0
Monounsaturated fat	18.0
Linoleic acid (omega-6 fatty acid)	16.0
$\alpha$ -Linolenic acid (omega-3 fatty acid)	57.0

(Source: Morris, D. H. 2003. Flax: A health and nutrition primer. p.11. Winnipeg: Flax Council of Canada)

oil. Ranhotra *et al.* (1992) noted that flaxseed oil or blends of flaxseed oil and sunflower oil promoted cholesterol reduction in hypercholesterolemic rats compared to diets formulated with hard fats. These authors suggested that a diet with the appropriate balance of n-6 and n-3 fatty acids was preferred over diets high in n-6 fatty acids. Ground flaxseed is high in omega-3 fatty acids which have been shown to reduce hypertension, cholesterol and triglyceride level (Oomah and Maza, 1998). Oikarinen *et al.* (2005) reported that flaxseed oil may be responsible for preventing colon carcinogenesis in multiple intestinal neoplasia (Min) mice. Dwivedi *et al.* (2005) also supported this finding that flaxseed oil prevented colon tumor development in rats. Presence of ALA in breast adipose tissue was inversely related to breast cancer risk (Maillard *et al.*, 2002). ALA, being the essential fatty acid, requirement can be fulfilled by intake of flaxseed products (Morris, 2004).

#### *Flaxseed as a source of protein*

The protein content in flaxseed has been reported to between 10.5% and 31% (Oomah and Mazza, 1993). Khategaon cultivars grown in India had a protein content of 21.9% (Madhusudhan and Singh, 1983). Differences in protein can be attributed to both genetics and environment. The proximate protein content of dehulled and defatted flaxseed varied considerably depending upon cultivar growth location and seed processing. Hull fraction contains lower protein levels

Table 3. Dietary Fiber content of flaxseed

Dietary Fiber component	Gram per 100 gram of flaxseed
Total Dietary	40
Soluble fiber	10
Insoluble fiber	30

(Source: Carter, J.F. 1993. Potential of flaxseed and flaxseed oil in baked goods and other products in human nutrition. *Cereal Foods World* 38 (10): 753-759)

and that dehulling increases protein level of flaxseed protein level from 19.2% to 21.8% (Oomah and Mazza, 1997). Albumin and globulin type proteins are the major proteins in flaxseed. Flaxseed albumin comprised 20% of meal protein (Madhusudhan and Singh, 1983). Globulin fraction makes up to 73.4% and the albumin constitutes about 26.6% of total protein (Marcone *et al.*, 1998). Flaxseed proteins are relatively high in arginine, aspartic acid and glutamic acid whereas lysine, methionine and cystine are limiting amino acid. Total amino acid content of the flaxseed after 8 days germination increased by 15 times with greatest increase (i.e. 200 times) being observed in glutamine and leucine compared to the original seed (Wanasundara *et al.*, 1999). Oomah and Mazza (1995) compared the nutritional value of flaxseed meal with soybean meal and concluded that net protein utilization and protein efficiency ratio of flaxseed meal were slightly lower than soybean meal with the exception of protein scores, which were high in flaxseed meal. The BV of flaxseed protein was similar to those of soybean protein (Frank, 1987). El-Kady (2000) found that the biological values of Belinka and Sakha-1 flaxseed proteins were 67.70 and 66.43 respectively.

Flaxseed protein was effective in lowering plasma cholesterol and triglycerides (TAG) compared to soy protein and casein protein (Bhathena *et al.*, 2002). Protein content of biscuits made from composite flour containing 15% ground flaxseed increased from 6.5% to 8.52%. The supplementation of flaxseed flour upto 15% showed no deleterious effect on the sensory attributes of biscuits (Zaib-un-Nisa, 2000). As flax is gluten-free, people who are sensitive to gluten can enjoy flax in their diets (Morris, 2003).

#### *Flaxseed as a source of dietary fiber (Mucilage or Gum)*

Dietary fiber is a communal word used to describe a variety of plant substances that are not easily digested by the enzymes responsible for digestion in humans (Eastwood and Passmore, 1983). Diets rich in dietary fibre may help reduce the risk of heart disease, diabetes, colorectal cancer, obesity and inflammation (Morris, 2003). Flaxseed is a rich source of dietary fiber (accounting 28% shown in Table 1), both soluble as well as insoluble fibers. Total dietary fiber content

of flaxseed is given in Table 3.

High amount of dietary fiber adds bulk to waste products in the gut and increases bile movement in the gastrointestinal movement. It exhibits natural laxative effect of dietary fiber. Flaxseed mucilage associated with hull of flaxseed is a gum like material composed of acidic and neutral polysaccharides. The neutral fraction of flaxseed contains xylose (62.8%) where as the acidic fraction of flaxseed is comprised mainly of rhamnose (54.5%) followed by galactose (23.4%) (Cui *et al.*, 1994). Low glycemic index foods containing soluble fiber not only prevent certain metabolic ramifications of insulin resistance but also reduce insulin resistance (Reaven *et al.*, 1993). Soluble fiber and other components of flaxseed fraction could potentially affect insulin secretion and its mechanism of action in maintaining plasma glucose homeostasis. Flaxseed was shown to reduce the post prandial blood glucose response in humans. Healthy female volunteers consumed 50 g ground, raw flaxseed/day for 4 weeks which provided 12-13% of energy intake (24-25 g/100 g total fat). Flaxseed raised  $\alpha$ -linolenic acid and long-chain  $\eta$ -3 fatty acids in both plasma and erythrocyte lipids, as well as raising urinary thiocyanate excretion 2.2 fold. Flaxseed also lowered serum total cholesterol by 9 % and low-density-lipoprotein-cholesterol by 18 %. Changes in plasma  $\alpha$ -linolenic acid were equivalent when 12 g  $\alpha$ -linolenic acid/day was provided as raw flaxseed flour (50 g/day) or flaxseed oil (20 g/day) suggesting high bioavailability of  $\alpha$ -linolenic acid from ground flaxseed. Test meals containing 50 g carbohydrate from flaxseed or 25 g flaxseed mucilage each significantly decreased postprandial blood glucose responses by 27 %. 50 g high- $\alpha$ -linolenic acid flaxseed/day is palatable, safe and may be nutritionally beneficial in humans by raising n-3 fatty acids in plasma and erythrocytes and by decreasing postprandial glucose responses (Cunnae *et al.*, 1993).

Similar findings were observed in post menopausal women fed 40 g/day flaxseed fortification diet (Lemay *et al.*, 2002). Bread containing 25% flaxseed gave a glycemic response that was 28% lower than the control (no flaxseed) bread (Jenkins *et al.*, 1999).

#### *Flaxseed as a source of lignan (Phenolic compounds)*

Flaxseed is the richest source of plant lignans (Thompson *et al.*, 1991). Secoisolariciresinol diglucoside (SDG) is the predominant lignan in flaxseed with minor amount of pinoresinol and matairesinol (MAT) (Meagher *et al.*, 1999; Thompson *et al.*, 1991). SDG was found 2653 mg/100 g of non defatted flaxseed extract (Hall and Shultz, 2001). The

lignans of flaxseed are phytoestrogens and serves as precursors in the production of mammalian lignans. Flaxseed lignans convert to mammalian lignans enterolactone and enterodiol by intestinal flora (Wang *et al.*, 2000). Kitts *et al.* (1999) reported that enterolactone and enterodiol had greater antioxidant activity than the present lignan (SDG), suggesting that the metabolites might be the reason for the health benefits of plant lignans. Prasad (2005) suggested that lignan may act to prevent oxygen radical production, thus effectively reducing atherosclerosis. Lignans have antioxidant activity and thus may contribute to the anticancer activity of flaxseed (Yuan *et al.*, 1999; Kangas *et al.*, 2002; Prasad 1997). However, number of factors may contribute to the various anticancer activity of flaxseed (Thompson *et al.*, 2005). The behavior of the lignans depends upon the biological level of estradiol. At normal estradiol levels, the lignans act as estrogen antagonists but in post menopausal women (i.e. low estradiol levels) can act as weak estrogen (Hutchins and Slavin, 2003; Rickard and Thompson, 1997). Although lignans have been shown to be protective against breast cancer, minor structural alterations may influence overall activity (Sarrinen *et al.*, 2005). Further research is needed to explore the role of flaxseed in cancer prevention. Toure and Xueming (2010) reviewed on flaxseed lignan which provides better understanding of flaxseed antioxidant activities and suggests that flaxseed lignans may be used as natural antioxidants.

Flaxseed, in particular lignans could influence bone development. Ward *et al.* (2001b) found that rats exposed to 88 or 177.3 mg SDG/kg of body weight/day had higher bone strength than the basal diet at 50 days post natal. However, by post natal day 132, no differences in bone strength, bone mineral density were observed. Exposure to SDG did not have negative effect on bone strength (Ward *et al.*, 2001b).

#### *Anti-nutrients in flaxseed*

Keeping an eye on safety of flaxseed, two compounds, cyanogenic glycosides and linatine an antipyridoxine factor are questioned frequently. Release of hydrogen cyanide from flaxseed would be minimal and below toxic lethal dose. At the recommend daily intake of about 1–2 table spoons, approximately 5–10 mg of hydrogen cyanide is released from flaxseed, which is well below the estimated acute toxic dose for an adult of 50–60 mg inorganic cyanide and below the 30–100 mg/day humans can detoxify (Roseling, 1994). Generally roasting is carried out to eliminate cyanogenic glycosides. Wanasundara *et al.* (1993) studied on

removal of cyanogenic glycosides of flaxseed meal by a two phase solvent extraction system consisting of hexanes and an alkanol (Methanol, ethanol or isopropanol) phase with or without added water and/or ammonia. Of the 4.42 mg/g linustatin and 1.90 mg/g neolinustatin originally present in the meals, over 90% of each cyanogenic glycoside was removed under optimum conditions using methanolic solutions. Although linatine is a problem in chicks, flaxseed has not been associated with a vitamin B6 deficiency in human. In fact, no affect on serum pyridoxine levels in subjects consuming 45 g/day of flaxseed over 5 weeks was observed (Dieken, 1992).

In addition to this, trypsin inhibitor and phytic acid are other antinutrients contained in flaxseed. But compared to soyabean and canola seeds, activity of them are low (Hall *et al.*, 2006). Bhatti (1993) reported laboratory-prepared flaxseed meals containing 42–51 units of TIA (Trypsin inhibitor activity), which was slightly higher than 10–30 units observed by Madhusudhan and Singh (1983) and commercially obtained flaxseed meal (14–37 units). The contents of phytic acid were significantly different among cultivars. AC Linora has a lowest phytic acid content of 2280 mg/100 g and low ALA yellow-seeded cultivar Linola 947 has the highest content (3250 mg/100 g seed) among the eight cultivars reported (Oomah *et al.*, 1996).

#### **Conclusion**

Flaxseed (Linseed) encompasses the potential health suiting nutritional profile in it. However, many people are still unaware of the potential health benefits of flaxseed and food applications. ALA (omega-3 fatty acid), dietary fiber and Lignan (specifically SDG) content attracts food technologists to explore its abilities at fullest extent in commercial food processing sector. Recently Baking and Pasta companies have incorporated flaxseed into their formulations. General recommendation for daily intake has been 1–3 table spoons per day for ground flaxseed or 1 table spoon for flaxseed oil. Flaxseed is emerging as one of the nutritive and functional ingredient in food products. Scientific findings are growing in support of flaxseed consumption. More studies are needed to resolve the conflicting reports regarding the health benefits, in particular the role of ALA and SDG in prostate cancer and cancer in general.

#### **References**



- Anonymous. 2000. Oil World Statistics Update. Oil World 31: 9-10.
- Berglund, D.R. 2002. Flax: New uses and demands. In Janick, J. and Whipkey, A. (Eds.). Trends in new crops and new uses, p. 358-360. Alexandria: ASHS Press.
- Bhathena, S.J., Ali, A.A., Mohamed, A.I., Hansen, C.T. and Velasquez, M.T. 2002. Differential effects of dietary flaxseed protein and soy protein on plasma triglyceride and uric acid levels in animal models. Journal of Nutritional Biochemistry 13 (11): 684-689.
- Bhatty, R.S. 1993. Further compositional analyses of flax: Mucilage, trypsin inhibitors and hydrocyanic acid. Journal of American Oil Chemist Society 70 (9): 899-904.
- Carter, J.F. 1993. Potential of flaxseed and flaxseed oil in baked goods and other products in human nutrition. Cereal Food World 38 (10): 753-75.
- Carter, J.F. 1996. Sensory evaluation of flaxseed of different varieties. In Proceedings of the 56<sup>th</sup> Flax Institute of the United States, p. 201-203. Fargo North Dakota: Flax Institute of United States.
- Chan, J.K., McDonald, B.E., Gerrad, J.M., Bruce, V.M., Weaver, B. J. and Holub, B.J. 1993. Effect of dietary alpha-linolenic acid and its ratio to linoleic acid on platelet and plasma fatty acids and thrombogenesis. Lipids 28 (9): 811-817.
- Clifford, H., Mehmetc, T. and Yingying, X. 2006. Flaxseed. In Taylor, S. L. (Ed). Advances in Food and Nutrition Research 51, p.1-97. Netherland: Elsevier Science and Technology, Academic Press.
- Cui, W., Mazza, G. and Biliaderis, C.G. 1994. Chemical structure, molecular size distributions and rheological properties of flaxseed gum. Journal of Agricultural and Food Chemistry 42 (9): 1891-1895.
- Cunnane, S.C., Ganguli, S. and Menard, C. 1993. High  $\alpha$ -linolenic acid flaxseed (*Linum usitatissimum*): Some nutritional properties in humans. British Journal of Nutrition 69 (2): 443-453.
- Daun, J.K., Barhet, V.J., Chornick, T.L. and Duguid, S. 2003. Structure, composition, and variety development of flaxseed. In Thompson, L. U. and Cunnane, S. C. (Eds). Flaxseed in Human Nutrition, 2nd ed, p. 1-40. Champaign, Illinois: AOCS Press
- Daun, J.K. and DeClercq, D.R. 1994. Sixty years of Canadian flaxseed quality surveys at the Grain Research Laboratory. In 55<sup>th</sup> Proceedings of Flax Institute of United States, p. 192-200. Fargo North Dakota: Flax Institute of United States.
- Diederichsen, A. and Richards, K. 2003. Cultivated flax and the genus *Linum* L.: taxonomy and gerplasm conservation. In Muir, A. D. and Westcott, N. D. (Eds). Flax, The genus *Linum*, p. 22-54. London: Taylor & Francis.
- Dieken, H.A. 1992. Use of flaxseed as a source of omega-3 fatty acids in human nutrition. In The 54<sup>th</sup> Proceeding of Flax Institute of United States. p. 1-4. Fargo North Dakota: Flax Institute of United States.
- Dwivedi, C., Natarajan, K. and Matthees, D. 2005. Chemopreventive effects of dietary flaxseed oil on colon tumor development. Nutrition and Cancer 51 (1): 52-58.
- Eastwood, M.A. and Passmore, R. 1983. Dietary fiber. The Lancet. 2 (8343): 202-206.
- El-Kady, E.A. 2000. Chemical and Technological studies on seeds of some filed crops. Faculty of Agriculture. Egypt: Kafr El-Sheikh, Tanta University, Ph.D. Thesis.
- Frank, A.W. (1987). Food Uses of Cottonseed protein. In Hudson, B. J. (Ed). Development in Food Proteins-5. p. 30-80. London: Elsevier Applied Science Publishers.
- Freeman, T.P. 1995. Structure of flaxseed. In Cunnane, S. C. and Thompson, L. U. (Eds). Flaxseed in Human Nutrition, p. 11-21. Champaign Illinois: AOCS Press.
- Hall, C. and Shultz, K. 2001. Phenolic antioxidant interactions. In Abstracts of the 92nd American Oil Chemists Society Annual Meeting and Expo. p. S88 Minneapolis, Minnesota: AOCS
- Hurteau, M.C. 2004. Unique new food products contain good omega fats. Journal of Food Science Education 3 (4): 52-53.
- Hutchins, A.M. and Slavin, J.L. 2003. Effect of flaxseed on sex hormone metabolism. In Thompson, L.U. and Cunnane, S.C. (Eds). Flaxseed in Human Nutrition. 2<sup>nd</sup> ed, p. 126-149. Champaign, Illinois: AOCS Press.
- Institute of Medicine. 2002. Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids. p. 8-97. Washington, DC: Food and Nutrition Board.
- Jenkins, D.J.A., Kendall, C.W.C. and Vidgen, E. 1999. Health aspects of partially defatted flaxseed, including effects on serum lipids, oxidative measures, and *ex vivo* androgen and progesterin activity: A controlled crossover trial. American Journal of Clinical Nutrition 69 (3): 395-402.
- Kangas, L., Saarinen, N. and Mutanen, M. 2002. Antioxidant and antitumor effects of hydroxymatairesinol (HM-3000, HMR), a lignan isolated from the knots of spruce. European Journal of Cancer Prevention 11 (2): S48-S57.
- Kitts, D.D., Yuan, Y.V., Wijewickreme, A.N. and Thompson, L.U. 1999. Antioxidant activity of the flaxseed lignan secoisolariciresinol diglycoside and its mammalian lignan metabolites enterodiol and enterolactone. Molecular and Cellular Biochemistry 202 (1-2): 91-100.
- Lemay, A., Dodin, S. and Kadri, N. 2002. Flaxseed dietary supplement versus hormone replacement therapy in hypercholesterolemic menopausal women. Obstetrics and Gynecology 100 (3): 495-504.
- Madusudhan, K.T. and Singh, N. 1983. Studies on linseed proteins. Journal of Agricultural and Food Chemistry 31 (5): 959-963.
- Maillard, V., Bougnoux, P. and Ferrari, P. 2002. Omega 3 and Omega 6 fatty acids in breast adipose tissue and relative risk of breast cancer in a case-control study in Tours, France. International Journal of Cancer 98 (1): 78-83.
- Marcone, M.F., Kakuda, Y. and Yada, R.Y. 1998. Salt-soluble seed globulins of dicotyledonous and monocotyledonous

- plants. II. Structural characterization. Food Chemistry 63 (2): 265–274.
- Meagher, L.P., Beecher, G.R., Flanagan, V.P. and Li, B.W. 1999. Isolation and characterization of the lignans, isolariciresinol and pinoresinol, in flaxseed meal. Journal of Agricultural Food Chemistry 47 (8): 3173–3180.
- Morris, D.H. 2003. Flax: A health and nutrition primer. 3<sup>rd</sup> ed, p.11 Winnipeg: Flax Council of Canada. Downloaded from <http://www.jitinc.com/flax/brochure02.pdf> verified on 4/6/12.
- Morris, D.H. 2004. Other health benefits of flax. In Flax: A Health and Nutrition Primer, Flax Council of Canada. Downloaded from [http://www.flaxcouncil.ca/english/pdf/FlxPrmr\\_4ed\\_Chpt7.pdf](http://www.flaxcouncil.ca/english/pdf/FlxPrmr_4ed_Chpt7.pdf) on 4/6/2012
- Morris, D.H. 2007. Flax - A Health and Nutrition Primer. 4<sup>th</sup> ed. Downloaded from <http://www.flaxcouncil.ca/english/index.jsp?p=primer&mp=nutrition> on 4/6/2012.
- Nettleton, J. 2003. Collected Recommendations for LC-PUFA Intake. PUFA Newsletter. Downloaded from <http://www.fatsoflife.com/wp-content/uploads/pdfs/PUFA0903.pdf> on 4/6/2012
- Oikarinen, S., Pajari, A., Salminen, I., Heinonen, S.M., Adlercreutz, H. and Mutanen, M. 2005. Effects of a flaxseed mixture and plant oils rich in alpha-linolenic acid on the adenoma formation in multiple intestinal neoplasia (Min) mice. British Journal of Nutrition 94 (4): 510–518.
- Oomah, B.D. 2001. Flaxseed as a functional food source. Journal of the Science of Food and Agricultural. 81 (9): 889-894.
- Oomah, B.D. and Mazza, G. 1998. Flaxseed products for disease prevention. In Mazza, G. (Ed). Functional Foods: Biochemical and Processing Aspects. p. 91-138. Lancaster, PA: Technomic Publication Company Inc.
- Oomah, B.D. and Mazza, G. 1997. Effect of dehulling on chemical composition and physical properties of flaxseed. Lebensmittel-Wissenschaft und-Technologie 30 (2): 135–140.
- Oomah, B.D. and Mazza, G. 1995. Functional properties and uses of flaxseed protein. Inform 6 (11): 1246-1252.
- Oomah, B.D. and Mazza, G. 1993. Flaxseed proteins—a review. Food Chemistry 48 (2): 109–114
- Prasad, K. 1997. Hydroxyl radical-scavenging property of secoisolariciresinol diglucoside (SDG) isolated from flax-seed. Molecular Cellular Biochemistry 168 (1-2): 117–123.
- Prasad, K. 2005. Hypocholesterolemic and antiatherosclerotic effect of flax lignan complex isolated from flaxseed. Atherosclerosis 179 (2): 269–275.
- Ranhotra, G.S., Gelroth, J.A. and Glaser, B.K. 1992. Lipidemic response to rats fed flaxseed or sunflower oils. Cereal Chemistry 69 (6): 623–625.
- Reaven, G.M., Brand, R.J., Chen, Y.D.I., Mathur, A.K. and Goldfine, I. 1993. Insulin resistance and insulin secretion are determinants of oral glucose tolerance in normal individuals. Diabetes 42 (9): 1324–1332.
- Rickard, S.E. and Thompson, L.U. 1997. Phytoestrogens and lignans: Effects on reproduction and chronic disease. In Shahidi, F. (Ed). Antinutrients and Phytochemicals in Foods. p. 273-293. New York: Oxford University Press
- Roseling, H. 1994. Measuring effects in humans of dietary cyanide exposure to sublethal cyanogens from Cassava in Africa. Acta Horticulturae No.375 : 271–283.
- Saarinen, N.M., Makela, S. and Satti, R. 2003. Mechanism of anticancer effects of lignans with a special emphasis on breast cancer. In Cunnane, S. C. and Thompson, L. U. (Eds). Flaxseed in Human Nutrition. 2<sup>nd</sup> ed, p. 223-228. Champaign, Illinois : AOCS Press
- Thompson, L.U., Chen, J.M., Li, T., Strasser-Weippl, K. and Goss, P.E. 2005. Dietary flaxseed alters tumor biological markers in postmenopausal breast cancer. Clinical Cancer Research 11 (10): 3828–3835.
- Thompson, L.U. 1995. Flaxseed lignan and cancer. In Cunnane, S. C. and Thompson, L. U. (Eds). Flaxseed in human nutrition. 2<sup>nd</sup> ed, p. 219-236. Champaign, Illinois: AOCS Press.
- Thompson, L.U., Robb, P., Serraino, M. and Cheung, F. 1991. Mammalian lignan production from various foods. Nutrition and Cancer 16 (1): 43–52.
- Toure, A. and Xueming, X. 2010. Flaxseed lignans: Source, biosynthesis, metabolism, antioxidant activity, bioactive components and health benefits. Comprehensive Reviews in Food Science and Food Safety. 9 (3): 261-269.
- Vaisey-Genser, M. and Morris, D.H. 2003. Introduction: history of the cultivation and uses of flaxseed. In Muir, A. D. and Westcott, N. D. (Eds). Flax: The genus *Linum*. p. 1-2. London: Taylor & Francis
- Wanasundara, P.K.J.P.D., Amarowicz, R., Karab, M.T. and Shahidi, F. 1993. Removal of cyanogenic glycosides of flaxseed meal. Food Chemistry. 48 (3): 263-266.
- Wanasundara, P.K.J.P.D., Wanasundara, U.N. and Shahidi, F. 1999. Changes in flax (*Linum usitatissimum* L.) seed lipids during germination. Journal of American Oil Chemists' Society 76 (1): 41–48.
- Wang, J., Rosell, C.M., Benedito, D.E. and Barber, C. 2002. Effect of the addition of different fibers on wheat dough performance and bread Quality. Food Chemistry 79 (2): 221-226.
- Ward, W.E., Yuan, Y.V., Cheung, A.M. and Thompson, L.U. 2001a. Exposure to flaxseed and its purified lignan reduces bone strength in young but not older male rats. Journal of Toxicology, Environment and Health 63 (Part A): 53–65.
- Ward, W.E., Yuan, Y.V., Cheung, A.M. and Thompson, L.U. 2001b. Exposure to purified lignan from flaxseed (*Linum usitatissimum*) alters bone development in female rats. British Journal of Nutrition 86 (4): 499–505.
- WHO/FAO Expert Consultation on Diet. 2003. Diet, nutrition and the prevention of chronic diseases. World Health Organization Technical Report Series 916. p. 89–90. Geneva: World Health Organization.
- Yuan, Y.V., Rickard, S.E. and Thompson, L.U. 1999.

Short-term feeding of flaxseed or its lignan has minor influence on *in vivo* hepatic antioxidants status in young rats. Nutrition Research 19 (8): 1233–1243.

Zaib-un-Nisa. 2000. Effect of flaxseed supplementation on chemical properties of biscuits. Faisalabad, Pakistan: Agriculture University, M.Sc. Thesis.