

Evaluation of antioxidant, anti-tyrosinase potentials and phytochemical composition of four Egyptian plants

*¹Rashed, K., ²Medda, R., ²Spano, D. and ²Pintus, F.

¹Department of Pharmacognosy, Pharmacy and Pharmaceutical Industries Division,
National Research Centre, 33-El-Bohouth st. (former El-Tahrir st.)-Dokki,
Giza, Egypt. P.O. 12622

²Department of Sciences of Life and Environment, University of Cagliari, Cittadella Universitaria,
I-09042 Monserrato, CA, Italy

Article history

Received: 19 November 2014

Received in revised form:

17 April 2015

Accepted: 11 June 2015

Abstract

The aim of this present research is to evaluate antioxidant and antityrosinase potentials and to investigate the phytoconstituents of methanol (70%) extract from four Egyptian plants, *Solanum rantonnetii*, *Tilia cordata*, *Cichorium intybus* (L.), and *Lagerstroemia tomentosa*. Antioxidant activity was measured by using two different free radical scavenging methods, 2,2'-Azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) and 2,2-diphenyl-1-picrylhydrazyl radical (DPPH•), whereas polyphenols and flavonoids contents were evaluated by using Folin-Ciocalteu, and aluminum nitrate methods respectively. Tyrosinase inhibition was evaluated using commercial enzyme and dihydroxyphenylalanine (DOPA) as substrate and methanol extract of each plant was investigated by phytochemical analysis. The results showed that all the plants had good antioxidant activity where *Cichorium intybus* exhibited the highest antioxidant activity determined as total content of free-radical scavenging, polyphenol and flavonoid molecules, and showed the best tyrosinase inhibition activity compared with the other plants extracts studied. Chromatographic separation and NMR analysis of *Cichorium intybus* methanol extract revealed the identification of two coumarins, scopoletin and esculetin, and seven flavonoids, dihydroquercetin 7-4'-dimethyl ether, blumeatin, diosmetin, tamarixetin, quercetin, quercetin 3-O-β-galactoside and kaempferol 3-O-rutinoside. These findings suggest that *Cichorium intybus* is rich with bioactive compounds and could be used as a good source of potentially natural antioxidants and antityrosinase molecules.

© All Rights Reserved

Keywords

Medicinal plants
Cichorium intybus
Aerial parts
Phytoconstituents
Antioxidants
Tyrosinase inhibition

Introduction

The medicinal plants represent an interesting contribution to modern therapeutics due to the presence in their composition of molecules with pharmacological and antioxidant action. Research on natural antioxidants has become increasingly active in various fields due to the action of antioxidant molecules on protection from the damage induced by oxidative stress. It is well known that free radicals may be involved in a number of diseases including cancer, cardiovascular diseases, diabetes, Alzheimer's disease, Parkinson's disease. Thus, many studies have been carried on antioxidants in order to identify some bioactive compounds from natural sources (Pintus *et al.*, 2013). Plant materials could be also a great source of powerful skin-whitening agents in medicinal and cosmetic fields (Chang, 2009). Tyrosinase (EC 1.14.18.1) is a copper-containing enzyme that catalyzes the first steps of melanin synthesis: hydroxylation of tyrosine and the oxidation of 3, 4-dihydroxyphenylalanine (L-DOPA)

to o-dopaquinone. Abnormal production of melanin is the cause of various dermatological disorders such as melasma, age spots and post-inflammatory hyperpigmentation. Because of their abilities to decrease melanin production, there is a big demand for tyrosinase inhibitors in order to threat dermatological disorders associated with skin hyperpigmentation. Four Egyptian medicinal plants including *Solanum rantonnetii*, *Tilia cordata*, *Cichorium intybus*, and *Lagerstroemia tomentosa* have been selected for the present study on the basis of their antioxidant and tyrosinase inhibitory properties. *Solanum* is a large and diverse genus of flowering plants, including two food crops of the highest economic importance, the potato and the tomato. *Solanum rantonnetii* is a shrub from Solanaceae family. The roots have soothing effects of the plant upon ingestion (Quattrocchi, 2000). *Tilia cordata* is a deciduous tree from Tiliaceae family. Active ingredients in the *Tilia* flowers include flavonoids (which act as antioxidants) and volatile oils. The plant also contains tannins that can act as

*Corresponding author.

Email: khalednabih2015@yahoo.co.uk

Tel: 01003642233

an astringent (Bradley, 1992). Infusions obtained from the flowers of *Tilia cordata* are widely used for the treatment of fever and anxiety problems in folk medicine. Recently some research have been done about the antioxidant capacity of flower extracts of this plant (Vinha *et al.*, 2013) but nothing it's known about the other aerial parts of the plant, object of this research. *Lagerstroemia tomentosa* from Lythraceae family is a medicinal and ornamental plant. The bark is considered as stimulant and febrifuge, leaves and flowers are used as purgative (Chopra *et al.*, 1958), and the roots are astringent (Kirtikar and Basu, 1935). *Cichorium intybus* (L.) known as chicory is a plant from Asteraceae family. This plant is native to Europe and Asia. It has been widely used in folk medicine for treatment of gallstones, appetite loss, gout, jaundice, skin swellings, rheumatism, and liver inflammation (Gazzani *et al.*, 2000). Historically, chicory was grown by the ancient Egyptians as a medicinal plant, coffee substitute, and vegetable crop and was occasionally used for animal forage. Important phytochemicals are distributed throughout the plant, but the main contents are present in the root (Bais and Ravishankar, 2001). The seeds extract of *C. intybus* has high antioxidant and antidiabetic activities (Ghamarian *et al.*, 2012), also it has ameliorating effects on non-alcoholic fatty liver disease (Ziamajidi *et al.*, 2013). Methanolic extract of *C. intybus* and its various fractions have revealed wound healing effects. The methanolic extract of the plant has demonstrated some anticancer and apoptosis inducing effects (Suntar *et al.*, 2012). The objective of this research is to explore phytoconstituents and potential activities of four Egyptian plants to contribute to the knowledge of these plants that might represent a potential source of bioactive natural molecules.

Materials and Methods

Materials

All chemicals were obtained as pure commercial products and used without further purification. 2,2'-Azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS), aluminum nitrate, 2,2-diphenyl-1-picrylhydrazyl radical (DPPH•), Folin-Ciocalteu phenol reagent, gallic acid, quercetin, mushroom tyrosinase and L-3,4-dihydroxyphenylalanine (L-DOPA) were from Sigma Chemical Co (St. Louis, USA).

UV/VIS: Shimadzu UV-visible recording spectrophotometer model-UV 240 (NRC, Egypt), and Ultrospec 2100 spectrophotometer (Biochrom Ltd, Cambridge, England) using cells with a 1

cm path length. ¹H-NMR and ¹³C-NMR spectra were collected on Inova NB Varian instrument (Varian Scientific Instruments, Palo Alto, CA). MS (Finnigan MAT SSQ 7000, 70 ev). Silica gel (0.063-0.200 mm for column chromatography) and Sephadex LH-20 (Pharmacia Fine Chemicals). Thin layer chromatography (TLC) F254 plates. Solvent mixtures, BAW (n-butanol:acetic acid: water 4:1:5, (v/v/v), upper phase, 15% acetic acid). Paper Chromatography (PC) Whatman No.1 (Whatman Led. Maid Stone, Kent, England) sheets for qualitative detection of flavonoids and sugars.

Plants identification and collection

Aerial parts (leaves and stems) from *Solanum rantonnetii* were collected from Zoo garden, Giza, Egypt. *Tilia cordata* and *Cichorium intybus* aerial parts were collected from the Agricultural Research Centre, Giza, Egypt, while aerial parts of *Lagerstroemia tomentosa* were collected from Al-Zohiriya garden, Giza, Egypt in May 2012. All the plants were identified by Dr. Mohammed El-Gebaly, Department of Botany, National research centre (NRC) and by Mrs. Tereza Labib consultant of plant taxonomy at the ministry of agriculture and director of Orman botanical garden, Giza, Egypt.

Preparation of the plants extracts

Air dried aerial parts of *S. rantonnetii* (280 g), *T. cordata* (340 g) *C. intybus* (640 g) and *L. tomentosa* (480 g) were extracted with methanol (70%) several times at room temperature by maceration method. Each extract was concentrated under reduced pressure to give 22.5 g, 25.5 g, 32 g and 28 g, respectively and immediately dissolved in DMSO before every experiment in order to have a 1mg/ml solution. Each extract was phytochemically screened according to (Yadav and Agarwala, 2011) and the antioxidant capacity, polyphenols, and flavonoids content were determined. Moreover, the ability of these extracts to inhibit tyrosinase activity was also evaluated.

ABTS⁺ assay

This method (Re *et al.*, 1999) is based on the capacity of an antioxidant to scavenge the free radical ABTS⁺. The ABTS⁺ was generated by reacting 7 mM ABTS with 2.45 mM potassium persulfate (final concentration) in aqueous solution and the mixture was kept in the dark at room temperature for 24 h before use. The ABTS⁺ solution, which has an absorption maximum at 734 nm, was diluted in the appropriate solvent to obtain an A₇₃₄ of about 0.70. Extracted samples at different concentration (10 µL) were added to 1 mL of diluted ABTS⁺ solution and

mixed vigorously. After reaction at room temperature for 1 min, the absorbance at 734 nm was measured. The decrease in A_{734} was calculated and the results are presented as the ability of antioxidant to scavenge 50% of free radical ABTS $^{+}$ (IC₅₀).

DPPH[•] assay

The assay was carried out as previously described (Pintus *et al.*, 2013). DPPH[•] radical shows a strong absorption maximum at 515 nm and goes through a change of color after the transfer of a hydrogen atom from an antioxidant. An aliquot of extracts at different concentration (25 μ L) was added to 975 μ L of 6×10^{-5} M DPPH[•] solution. The reaction mixture was incubated in the dark for 15 min and, thereafter, the A_{515} was recorded against DPPH[•] solution alone as a control. The decrease in absorbance of DPPH[•], after addition of test extracts compared to the control, was used to calculate the antioxidant activity which was expressed as ability of antioxidant to scavenge 50% of DPPH radical (IC₅₀).

Determination of total polyphenol content

Total polyphenol content in the extracts were determined according to the Folin-Ciocalteu method using a calibration curve obtained with gallic acid as polyphenolic reference standard (Mansouri *et al.*, 2005). Different concentrations of the extracts (10 μ L) were mixed with 50 μ L of the Folin-Ciocalteu reagent and 790 μ L of distilled water. After 1 min, 150 μ L of saturated sodium carbonate (20% aqueous solution) was added to the mixture. The mixture was kept in the dark for 45 min, and then the A_{750} was measured and compared to the gallic acid calibration curve. The polyphenol content was expressed as gallic acid equivalents (GAE; mM).

Determination of total flavonoid content

Total flavonoid content was determined as reported (Moreno *et al.*, 2000). The extracts (0.5 mL) were added to 0.1 mL of 10% aluminum nitrate, 0.1 mL of 1 M sodium acetate and 4.3 mL of 80% EtOH. After 40 min at room temperature, the absorbance was determined at 415 nm. Flavonoid concentration was calculated by a calibration curve obtained using quercetin as reference standard and expressed as quercetin equivalent molarity (QE, mM).

Mushroom tyrosinase activity

The inhibitory effect of the plant extracts on tyrosinase activity was determined spectrophotometrically with the degree of inhibition of mushroom tyrosinase-catalysed oxidation of L-DOPA (Rahaman *et al.*, 2001). The reaction mixture

contained the extracts and L-DOPA (1.25 mM) in 25 mM phosphate buffer (pH 6.8). Then, tyrosinase (100 U/ml) was added into the mixture and the activity was determined by following the increase in absorbance at 475 nm resulting from the formation of the dopachrome product. The inhibition of tyrosinase activity was calculated with the following formula: Inhibition (%) = $(1 - (A_{475} \text{ in sample}/A_{475} \text{ in control})) \times 100\%$.

Isolation of the components of methanol (70%) extract from *Cichorium intybus* aerial parts

Methanol (70%) extract of *C. intybus* (28 g) was subjected to silica gel column chromatography eluting with n-hexane, dichloromethane, ethyl acetate and methanol gradually. One hundred and twenty fractions of 100 ml conical flask were collected. The fractions that showed similar paper chromatography in two solvent systems, butanol-acetic acid-water (BAW, (v/v/v) 4:1:5 and 15% acetic acid were combined to give four main fractions. Fraction 1 (4.55 g) was subjected to sub-column of silica gel eluted with n-hexane: dichloromethane where elution with n-hexane-dichloromethane (60:40) yielded compound 1 and further elution with n-hexane:dichloromethane (80:20) yielded compound 2. Fraction 2 (3.85 g) was subjected to sub-column of silica gel eluted with dichloromethane:ethyl acetate. Elution with dichloromethane: ethyl acetate (90:10) gave compound 3 and further elution with dichloromethane:ethyl acetate (85:15) yielded compound 4. Fraction 3 (6.45 g) was subjected to sub-column of silica gel eluted with dichloromethane: methanol. Elution with dichloromethane:methanol (98:2) afforded compound 5 while elution with dichloromethane:methanol (95:5) yielded compound 6 and further elution with dichloromethane:methanol (90:10) gave compound 7. Fraction 4 (5.65 g) was subjected to sub-column of silica gel eluted with ethyl acetate:methanol (95:5) afforded compound 8 and further elution with ethyl acetate:methanol (70:30) yielded compound 9. All the isolated compounds were purified on sephadex LH-20 column using methanol, ethyl alcohol and different systems of methanol and distilled water (methanol: distilled water, 1:1, 2:1, v/v).

General method for acid hydrolysis

Compounds 8, and 9 (5 mg) was heated in 5 ml 10% HCl for 5 h. The aglycones were extracted with ethyl acetate and identified by co-TLC with authentic standards. The sugars in the aqueous layer were identified by co-paper chromatography (co-PC) with authentic markers on Whatman No. 1 sheets in

Table 1. The radical scavenging activity of the four methanolic extracts determined with ABTS and DPPH assays and compared with Trolox, used as standard molecule

Methanol (70%) extract	IC ₅₀ values (µg/mL)	
	ABTS scavenging	DPPH scavenging
<i>S. rantonnetii</i>	57.9 ± 4.6	110.17 ± 7.6
<i>T. cordata</i>	139.9 ± 8.7	116.72 ± 12.8
<i>C. intybus</i>	21.56 ± 2.1	30.06 ± 2.6
<i>L. tomentosa</i>	57.64 ± 6.3	98.05 ± 7.8
Trolox	3.4 ± 0.4	3.2 ± 0.2

solvent system (n-BuOH-AcOH-H₂O 4:1:5 upper layer).

Results and Discussions

This study focused on antioxidant and antityrosinase activities of four methanol (70%) extracts of *Solanum rantonnetii*, *Tilia cordata*, *Cichorium intybus* and *Lagerstroemia tomentosa* and also phytochemical analysis was done.

The antioxidant ability of the extracts was screened by ABTS and DPPH assays and IC₅₀ values are reported in Table 1. *S. rantonnetii* and *L. tomentosa* showed the same scavenging capacity in each assay with an IC₅₀ of about 50 and 100 µg/mL for ABTS and DPPH methods, respectively. *T. cordata* has the lowest amount of antioxidant, whereas *C. intybus* showed the best antioxidant activity with lower IC₅₀ values of 21.56 and 30.06 µg/mL using ABTS, and DPPH methods.

These values are high if compared with trolox (IC₅₀~3.2 µg/mL); this is not surprising since analyzed samples are complex extracts and they most likely contain pro-oxidants agents which may compete with the antioxidants in the reaction with ABTS and DPPH radicals. Thus IC₅₀ values are the results of the balance between pro-oxidant and antioxidant molecules. In Figure 1A., the total polyphenolic content of the plants extracts is reported, expressed as gallic acid equivalent and measured by Folin-chocalteu method. Starting with the better extract, the amount of polyphenols follows the order *C. intybus* (2.33 mM) > *L. tomentosa* (1.02 mM) > *S. rantonnetii* (0.69 mM), and *T. cordata* (0.46 mM). The total flavonoid content also varies in the different extracts as shown in Figure 1B., *L. tomentosa* gave the highest value (0.5 mM) followed by *C. intybus* (0.25 mM) whereas *S. rantonnetii*, and *T. cordata* showed the lower values (0.113 mM, and 0.132 mM). Moreover, in vitro tyrosinase assay was performed to evaluate if

Table 2. Tyrosinase inhibition activity of the four methanolic plant extracts and IC₅₀ value of arbutin, used as standard molecule

Methanol (70%) extract	IC ₅₀ (mg/ml)
<i>S. rantonnetii</i>	-
<i>T. cordata</i>	0.67 ± 0.04
<i>C. intybus</i>	0.34 ± 0.03
<i>L. tomentosa</i>	-
Arbutin	0.13 ± 0.02

these plants extracts showed any inhibitory activity toward this enzyme. Melanin synthesis is carried out by a specific enzymatic pathway controlled by tyrosinase. The inhibition of tyrosinase is therefore the most common approach for the identification of new depigmenting agent. There is no any observed inhibitory effect for *S. rantonnetii*, and *L. tomentosa* extracts whereas, in presence of the extracts from *T. cordata* and *C. intybus*, the activity of mushroom tyrosinase was markedly decreased with the latest showing the strongest inhibitory effect (Table 2). Moreover, as compared with arbutin (IC₅₀ value of 0.13 mg/mL), a well known potent tyrosinase inhibitor, extracts from *T. cordata* and *C. intybus* had almost similar activity with IC₅₀ values of 0.67 and 0.34 mg/mL, respectively (Loizzo *et al.*, 2012). According to these results, our attention was focused on identification and characterization of the bioactive compounds present in *C. intybus* methanol extract. Phytochemical analysis of the four methanol (70%) extracts was done and the results are reported in Table 3, showing their phytoconstituents. *S. rantonnetii* extract proved the presence of triterpenes, flavonoids, tannins, and carbohydrates. *T. cordata* extract showed the presence of coumarins, triterpenes, flavonoids, tannins, saponins, and carbohydrates. *L. tomentosa* extract revealed the presence of triterpenes, flavonoids, tannins, and carbohydrates, while *C. intybus* extract showed that it contained triterpenes, flavonoids, tannins, carbohydrates, and coumarins (Table 3). Chromatographic separation of *C. intybus* methanol (70%) extract revealed the identification of two coumarins, scopoletin and esculetin, and seven flavonoids dihydroquercetin 7-4'-dimethyl ether, blumeatin, diosmetin, tamarixetin, quercetin, quercetin 3-O-β-galactoside and kaempferol 3-O-rutinoside (Figure 2). The NMR parameters for the identification of the isolated compounds are reported in the next paragraph

Structure elucidation of the isolated compounds

Scopoletin (1): 12 mg. ¹H-NMR (CDCl₃, 300




Figure 1. Polyphenol and flavonoid contents in the methanol (70%) extract of the four Egyptian plants. Panel A: Polyphenol amount is expressed as gallic acid equivalents (GAE); Panel B: The amount of flavonoids is expressed as quercetin equivalents (QE). All the extracts are prepared as 1mg/ml. The reported values are the mean of triplicate experiments

MHz) δ ppm 3.76 (3H, s, 6-OCH₃), 6.34 (1H, d, J=9.7 Hz, H-3), 7.08 (1H, s, H-5), 7.14 (1H, s, H-8), 7.72 (1H, d, J=9.7 Hz, H-4). ¹³C-NMR (CDCl₃, 100 MHz) δ ppm 56.34 (-OCH₃), 104.14 (C-8), 109.32 (C-5), 110.46 (C-10), 112.42 (C-3), 144.28 (C-4), 146.28 (C-6), 151.42 (C-7), 151.68 (C-9), 160.82 (C-2). It was obtained as colourless needles. NMR spectral data was very similar to that described by Kang *et al.*, (2003).

Esculetin (2): 10 mg. ¹H NMR (DMSO, 400 MHz): δ ppm 7.84 (1H, d, J=9.5 Hz, H-4), 6.98 (1H, s, H-5), 6.78 (1H, s, H-8), 6.14 (1H, d, J=9.5 Hz, H-3). ¹³C-NMR (DMSO, 100 MHz) δ ppm 160.84 (C-2), 150.48 (C-7), 149.12 (C-9), 144.58 (C-4), 143.28 (C-6), 112.85 (C-5), 111.86 (C-3), 116.38 (C-3), 110.86 (C-10), 103.16 (C-8). It was obtained as colourless needles and its spectral data was in agreement with that described by Lyi *et al.*, (2004).

Dihydroquercetin 7-4'-dimethyl ether (3): 18.5 mg. UV max (MeOH): 208, 232sh, 290, 336sh (NaOMe): 214, 248sh, 290, 356; (AlCl₃): 208, 224, 314, 384; (AlCl₃/HCl): 208, 224, 286sh, 312, 382sh;




Figure 2. Chemical structures of the compounds isolated from *C. intybus* aerial parts

(NaOAc): 229, 290, 358sh; (NaOAc/H₃BO₃): 232, 290, 336sh. ¹H-NMR (DMSO, 400 MHz): δ ppm 3.79 (3H, s, 4'-OCH₃); 3.82 (3H, s, 7-OCH₃), 4.58 (1H, dd, J=6.4, 11.4 Hz, H-3), 5.12 (1H, d, J=11.4 Hz, H-2), 5.88 (1H, d, J=6.4 Hz, 3-OH), 6.14 (1H, d, J=2.4, H-6), 6.18 (1H, d, J=2.4 Hz, H-8), 6.92 (1H, d, J=1.6 Hz, H-6'), 6.96 (1H, s, H-2'), 6.98 (1H, d, J=1.6 Hz, H-5'). ¹³C-NMR: (DMSO, 100 MHz): 197.85 (C4), 167.95 (C7), 163.15 (C5), 162.95 (C9), 148.76 (C4'), 146.38 (C3'), 129.66 (C1'), 119.48 (C6'), 115.24 (C2'), 112.35 (C5'), 101.58 (C10), 95.14 (C6), 94.26 (C8), 84.28 (C2), 71.76 (C3), 56.18 (7-OCH₃), 55.84 (4'-OCH₃). It was obtained as a white powder. Its spectral signals were in agreement with those described by Fazilatun *et al.*, (2004).

Blumeatin (4): 14.5 mg. UV max (MeOH): 208, 229sh, 288, 332sh; NaOMe): 214, 245sh, 288, 332sh; (AlCl₃): 212, 310, 378; (AlCl₃/HCl): 208, 224, 309, 375; (NaOAc): 230, 288, 332sh; (NaOAc/H₃BO₃): 232, 288, 332sh. ¹H-NMR (DMSO, 400 MHz): δ ppm 2.74 (1H, dd, J=2.9, 17.2 Hz, H-3a), 3.26 (1H, dd, J=12.6, 17.2 Hz, H-3b), 3.79 (3H, s, 7-OCH₃), 5.44 (1H, dd, J=2.9, 12.6 Hz, H-2), 6.08 (1H, d, J=2 Hz, H-6), 6.10 (1H, d, J=2 Hz, H-8), 6.78 (2H, s, H-2', H-6'), 6.89 (1H, s, H-4'); 12.15 (1H, s, 5-OH). ¹³C-NMR (DMSO, 100 MHz): δ

Table 3. Phytochemical analysis of the four methanolic plants aerial parts extracts

Phytoconstituents	<i>S. rantonnetii</i>	<i>T. cordata</i>	<i>C. intybus</i>	<i>L. tomentosa</i>
Triterpenes and /or Sterols	+	+	+	+
Carbohydrates and/or glycosides	+	+	+	+
Flavonoids	+	+	+	+
Coumarins	-	+	+	-
Alkaloids and/or nitrogenous compounds	-	-	-	-
Tannins	+	+	+	+
Saponins	-	+	-	-

(+) the presence of the constituents, (-) the absence of the constituents

ppm 197.15 (C4), 167.76 (C7), 163.38 (C5), 162.95 (C9), 145.9 (C3', C5'), 129.85 (C1'), 118.16 (C4'), 115.48 (C6'), 114.52 (C2'), 102.78 (C10), 94.75 (C6), 93.94 (C8), 78.82 (C2), 42.28 (C3), 56.45 (7-OCH₃). It was obtained as white powder. Its spectral signals were in agreement with those described by Fazilatun *et al.*, (2004).

Diosmetin (5): 16.5 mg. UV λ_{max} (MeOH): 247, 264, 345, (NaOMe): 249, 302, 388, (AlCl₃): 273, 352, 400 (AlCl₃/HCl): 272, 352, 400, (NaOAc): 252, 382, (NaOAc/H₃BO₃): 252, 382. 1H-NMR (DMSO-d₆, 500 MHz): δ ppm 7.65 (1H, d, J= 2.4 Hz, H2'); δ 7.35 (1H, dd, J=7.6, 2.4 Hz, H-6'); δ 6.85 (1H, d, J= 7.6 Hz, H-5'); δ 6.68 (1H, d, J=2.4 Hz, H-8); δ 6.45 (1H, d, J=2.2 Hz, H6); δ 6.65 (1H, s, H3); δ 3.9 (3H, s, OCH₃). 13C-NMR (DMSO-d₆, 100 MHz): δ ppm 162.9 (C-2), 103.8 (C-3), 182.6 (C-4), 161.4 (C-5), 98.7.0 (C-6), 164.7 (C-7), 94.2 (C-8), 157.6 (C-9), 104.2 (C-10), 123.4 (C-1'), 112.8 (C-2'), 147.3 (C-3'), 151.4 (C-4'), 112.7 (C-5'), 119.1 (C-6') and 55.6 (OCH₃). It was obtained as yellow powder. Its spectral data is in agreement with signals described by Lunesa *et al.*, (2011).

Tamarixetin (6): 12.5 mg. 1H NMR (DMSO-d₆, 300 MHz): δ ppm 7.84 (1H, d, J= 1.6 Hz, H-2'), 7.83 (1H, dd, J= 7.4 Hz, 1.6 Hz, H-6'), 6.98 (1H, d, J= 7.4 Hz, H-5'), 6.56 (1H, d, J= 1.9 Hz, H-8), 6.26 (1H, d, J= 1.9 Hz, H-6), 3.86 (3H, s, OMe). 13C-NMR (DMSO-d₆, 100 MHz): δ ppm 175.86 (C-4), 164.26 (C-7), 161.12 (C-5), 156.85 (C-9), 148.72 (C-4'), 146.68 (C-2), 146.62 (C-3'), 136.26 (C-3), 122.35 (C-10), 121.68 (C-6'), 115.54 (C-5'), 112.28 (C-2'), 103.65 (C-5), 98.25 (C-6), 93.64 (C-8). EI-MS: m/z 316. It was obtained as a yellow powder and it showed signals were very similar to that described by Benahmed *et al.*, (2011).

Quercetin (7): 6 mg. UV λ_{max} (MeOH): 255, 267, 371; (NaOMe): 270, 320, 420; (AlCl₃): 270, 455; (AlCl₃/HCl): 264, 303sh, 315sh, 428; (NaOAc): 257, 274, 318, 383; (NaOAc/H₃BO₃): 259, 387. EI-MS: m/z 302. It was obtained as yellow powder and

its spectral data is in agreement with Manguro *et al.*, (2005).

Quercetin 3-O- β -galactoside (8): 19 mg. UV λ_{max} (MeOH): 254, 266, 289, 352; (NaOMe): 275, 329, 406; (AlCl₃): 276, 304, 353, 435; (AlCl₃/HCl): 275, 296, 362, 404; (NaOAc): 276, 327, 390; (NaOAc/H₃BO₃): 266, 298, 372. 1H NMR (CD₃OD, 300 MHz): 6.14 (1H, d, J= 2 Hz, H-6), 6.34 (1H, d, J= 2 Hz, H-8), 6.87 (1H, d, J= 8.2 Hz, H-5'), 7.58 (1H, dd, J= 2, 7.7 Hz, H-6'), 7.84 (1H, d, J= 2 Hz, H-2'), 5.08 (1H, d, J= 7.8 Hz, H-1''), 3.9-3.1 (rest of sugar protons, H-2''-6''). It was obtained as yellow needles. Acidic hydrolysis gave quercetin as an aglycone and galactose as sugar and spectral signals were in accordance with that described by (Guvenalp and Demirezer, 2005).

Kaempferol 3-O-rutinoside (9): 24 mg. UV λ_{max} (nm): MeOH 268, 348; (NaOMe) 276, 326, 398; (AlCl₃): 275, 306, 357, 396; (AlCl₃/HCl) 275, 305, 354, 392sh; (NaOAc) 275, 314, 393; (NaOAc/H₃BO₃) 267, 356. 1H NMR (DMSO-d₆, 300 MHz), δ ppm 7.96 (2H, d, J= 8.5 Hz, H-2', 6'), 6.86 (2H, d, J= 8.5 Hz, H-3', 5'), 6.42 (1H, s, H-8), 6.24 (1H, s, H-6), 5.34 (1H, d, J= 7.2 Hz, H-1''), 4.42 (1H, s, H-1''), 3.0-4.1 (rest of rhamnose and glucose protons, 1.12 (3H, d, J= 6.5 Hz, -CH₃-rhamnosyl). 13C-NMR (DMSO-d₆, 100 MHz): δ ppm 156.44 (C-2), 133.15 (C-3), 177.28 (C-4), 161.85 (C-5), 98.75 (C-6), 164.45 (C-7), 93.85 (C-8), 156.74 (C-9), 103.78 (C-10), 120.75 (C-1'), 131.24 (C-2', 6'), 115.26 (C-3', 5'), 159.86 (C-4'), 101.38 (C-1''), 74.16 (C-2''), 76.35 (C-3''), 69.84 (C-4''), 75.62 (C-5''), 66.84 (C-6''), 100.75 (C-1'''), 70.25 (C-2'''), 70.52 (C-3'''), 71.74 (C-4'''), 68.15 (C-5'''), 17.62 (C-6'''). It was obtained as yellow amorphous powder. Acidic hydrolysis afforded kaempferol as an aglycone, rhamnose and glucose as sugar moieties. All data were identical with that described by Bao *et al.*, (2003). The results indicated that *C. intybus* exhibited the best antioxidant activity determined as total content of free-radical scavenging, and polyphenol molecules. Moreover, *C. intybus* extract

showed the highest value in terms of tyrosinase inhibitory activity. This observed activities can be explained by the properties of the isolated compounds and also phytoconstituents present in the extract, such as coumarins and flavonoids.

Plants contain a mixture of many kinds of secondary metabolism products, including phenols, which vary greatly in their antioxidant capacity. Coumarins are a large family of natural and synthetic compounds, which present different pharmacological activities. Recent studies paid special attention to their antioxidative, antiinflammatory, anticancer and enzymatic inhibition properties. In recent studies, some coumarins proved to be mushroom tyrosinase inhibitors (Borges *et al.*, 2009). As showed in Table 3, coumarins are absent in *S. rantonnetii* and *L. tomentosa* extracts but are present in the extracts from *T. cordata*, and *C. intybus*, which are the only extracts that showed tyrosinase inhibition activity. Thus, we could assume that coumarins can be responsible for the tyrosinase inhibitory capacity of *T. cordata* and *C. intybus* extracts. Flavonoids have anti-inflammatory, anti-allergic, anticarcinogenic, antioxidant, and antiviral properties (Cook, 1996). Also flavonoids provide potent protection against oxidative and free radical damage (Świgło and Tyrakowska, 2003). Several reports have suggested that compounds yielding antioxidant properties could also serve as potential antityrosinase agents capable of blocking melanin synthesis (Niki *et al.*, 2011). Compounds isolated from *C. intybus* methanol extract as dihydroquercetin 7-4'-dimethyl ether and blumeatin exhibited competitive inhibitor on tyrosinase activity (Saewan *et al.*, 2011). Dietary phenolics have been shown to possess antityrosinase and antioxidant properties as quercetin compound which was isolated from our plant extract exhibited antityrosinase and antioxidant effects (Weerawon *et al.*, 2014).

Conclusion

The results of this research indicated that methanol (70%) extract of aerial parts from four Egyptian plants possessed significant antioxidant activity determined as total content of free-radical scavenging, polyphenol, and flavonoid molecules and two of them showed a great tyrosinase inhibitory activity. Our results contribute to the knowledge of the components of these plants with a better characterization of the components of *C. intybus* extract which exhibits the best activities. The presence of antioxidant molecules and tyrosinase inhibitors make these plants a promising source of biomolecules

for applications in the pharmaceutical field. Many therapeutic options exist, though treatment is often difficult, requiring lengthy therapy and the use of these substances is often limited because of skin irritation caused by toxicity or occurrence of dermatitis. Thus, it has become increasingly necessary to search for new drugs. The next step will be the evaluation of antioxidant and antityrosinase activities of each isolated compound and the skin-whitening activity in cellular and *in vivo* systems.

References

- Bais, H.P. and Ravishankar, G.A. 2001. "Cichorium intybus L.—cultivation, processing, utility, value addition and biotechnology, with an emphasis on current status and future prospects," Journal of the Science of Food and Agriculture 81(5): 467–484.
- Bao, T.N., Peng, S.L. and Zhou, Z.Z. 2003. Chemical Constituents of Buckwheat. Natural Product Research and Development 15: 24-26.
- Benahmed, M., Akkal, S., Elomri, A., Laouer, H, Ve'rite, P. and Seguin, E. 2011. Constituents from *Bupleurum montanum* (Coss. and Dur.) (Apiaceae). Arabian Journal of Chemistry. doi:10.1016/j.arabjc.2011.01.001.
- Borges, F., Roleira, F., Milhazes, N., Uriarte, E. and Santana, L. 2009. Simple coumarins: privileged scaffolds in medicinal chemistry. Frontier Journal of Medicinal Chemistry 4: 23-85.
- Bradley, P., ed. (1992). British Herbal Compendium 1: 142–144.
- Chang, T.S. 2009. An Updated Review of Tyrosinase Inhibitors. International Journal of Molecular Sciences 10: 2440-2475.
- Chopra, R.N., Nayar, S.L. and Chopra, I.C. 1958. In Glossary & Indian Medicinal Plants, CSIR, New Delhi, 546-554.
- Cook, N.C. and Samman, S. 1996. Flavonoids-chemistry, metabolism, cardio protective effects, and dietary sources. Nutrition Biochemistry 7: 66- 76.
- Fazilatun, N., Zhari, I., Nornisah, M., Mas, R. and Hakim, M.H. 2004. Free radical-scavenging activity of organic extracts and of pure flavonoids of *Blumea balsamifera* DC leaves. Food Chemistry 88: 243–252.
- Gazzani, G., Daglia, M., Papetti, A. and Gregotti, C. 2000. *In vitro* and *ex vivo* anti- and prooxidant components of *Cichorium intybus*. Journal of Pharmaceutical and Biomedical Analysis 1: 127–133.
- Ghamarian, A., Abdollahi, M., Su, X., Amiri, A., Ahadi, A. and Nowrouzi, A. 2012. Effect of chicory seed extract on glucose tolerance test (GTT) and metabolic profile in early and late stage diabetic rats. DARU 15: 56-64.
- Guvenalp, Z. and Demirezer, L.O. 2005. Flavonol Glycosides from *Asperula arvensis* L. Turkish Journal of Chemistry 29: 163-169.
- Kang, W.Y., Li, G.-H. and Hao, X.-J. 2003. Two New Triterpenes from *Neonauclea sessilifolia*. Acta Botanica Sinica 45(8): 1003-1007.

- Kirtikar, K.R. and Basu, B.D. 1935. Indian Medicinal Plant Vol. III. International Book Distributors and Publishers, Dehradun, India, 71-89.
- Loizzo, M.R., Tundis, R. and Menichini, F. 2012. Natural and synthetic tyrosinase inhibitors as Antibrowning agents: an update. *Comprehensive Reviews in Food Science and Food Safety* 11: 378-398.
- Lunesa, C.P., Mylene, M.U., Kung, H.S., Mingfu, W. and Ivan K.C. 2011. Isolation and characterization of antimicrobial, anti-inflammatory and chemopreventive flavones from *Premna odorata* Blanco. *Journal of Medicinal Plants Research* 5(13): 2729- 2735.
- Lyi, L., Li-Wen, X., Yan-Feng, J., Chan-Juan, X., Han-Qing, W. and You-Rui, S. 2004. Isolation, Characterization and Crystal Structure of Natural Eremophilanolide from *Ligularia sagitta*. *Zeitschrift für Naturforschung* 59b: 921–924.
- Manguro, L.O.A., Ugi, I. and Lemmen, P. 2005. Flavonol Glycosides from the Leaves of *Embelia keniensis*. *Journal of the Chinese Chemical Society* 52: 201-208.
- Mansouri, A., Embarek, G., Kokkalou, E. and Kefalas, P. 2005. Phenolic profile and antioxidant activity of the Algerian ripe date palm fruit (*Phoenix dactylifera*). *Food Chemistry* 89: 411–420.
- Moreno, M.I.N., Isla, M., I., Sampietro, A.R. and Vattuone, M.A. 2000. Comparison of the free radical-scavenging activity of propolis from several regions of Argentina. *Journal of Ethnopharmacology* 71: 109–114
- Niki, Y., Yoshida, M., Ando, H., Wakamatsu, K., Ito, S. and Harada, N. 2011. 1-(2,4-Dihydroxyphenyl)-3-(2,4-dimethoxy-3-methylphenyl) propane inhibits melanin synthesis by dual mechanisms. *Journal of Dermatology Sciences* 63(2): 115-21.
- Pintus, F., Spanò, D., Mascia, C., Macone, A., Floris, G. and Medda, R. 2013. Acetylcholinesterase inhibitory and antioxidant properties of *Euphorbia characias* latex. *Records of Natural Products* 7: 147-151.
- Quattrocchi, U. 2000. *CRC World Dictionary of Plant Names*. USA: Taylor and Francis. p. 2058. ISBN 978-0-8493-2678-3.
- Rahaman, A.U., Choudhary, M.I. and Thomsen, W.J. 2001. Bioassay techniques for drug development. In: Thomsen WJ, editor. *Manual of Bioassay Techniques for Natural Products Research*. 1st edition. The Netherlands: Harwood Academic Publishers; p p 34–35.
- Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M. and Rice-Evans, C. 1999. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine* 26:1231–1237.
- Saewan, N., Koysomboon, S. and Chantrapromma, K. 2011. Anti-tyrosinase and anti-cancer activities of flavonoids from *Blumea balsamifera* DC. *Journal of Medicinal Plants Research* 5(6): 1018-1025.
- Suntar, I., Kupeli, A. E., Keles, H., Yesilada, E., Sarker, S.D. and Baykal, T. 2012. Comparative evaluation of traditional prescriptions from *Cichorium intybus* L. for wound healing: stepwise isolation of an active component by in vivo bioassay and its mode of activity. *Journal of Ethnopharmacology* 30: 299–309.
- Świąło, G. and Tyrakowska, B. 2003. Quality of Commercial Apple Juices Evaluated on the Basis of the Polyphenol Content and the TEAC Antioxidant Activity. *Journal of Food Sciences* 68(5): 1844-1849.
- Weerawon, T., Saowalak, L., Thanyawan, P. and Uraiwan, P. 2014. Comparative Evaluation of Antityrosinase and Antioxidant Activities of Dietary Phenolics and their Activities in Melanoma Cells Exposed to UVA. *Siriraj Medical Journal* 66(1): 5-10.
- Vinha, A.F., Sérgio, V.P.B., Ana, C. and Marisa, M. 2013. Comparison Between the Phytochemical and Antioxidant Properties of Plants Used in Plant Infusions for Medicinal Purposes. *Journal of Agricultural Science* 5(11): 11-19.
- Yadav, R.N.S. and Agarwala, M. 2011. Phytochemical analysis of some medicinal plants. *Journal of Phytology* 3(12): 10-14.
- Ziamajidi, N., Khaghani, S., Hassanzadeh, G., Vardasbi, S., Ahmadian, S., Nowrouzi, A., Ghaffari, S.M. and Abdirad, A. 2013. Amelioration by chicory seed extract of diabetes- and oleic acid-induced non-alcoholic fatty liver disease (NAFLD)/non-alcoholic steatohepatitis (NASH) via modulation of PPAR α and SREBP-1. *Food Chemistry and Toxicology* 58: 198–209.